



Cancer nanotechnology: The impact of passive and active targeting in the era of modern cancer biology[☆]



Nicolas Bertrand^{a,1}, Jun Wu^{b,1}, Xiaoyang Xu^{a,b}, Nazila Kamaly^b, Omid C. Farokhzad^{b,*}

^a The David H. Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^b Laboratory of Nanomedicine and Biomaterials, Department of Anesthesiology, Brigham and Women's Hospital, Harvard Medical School, 75 Francis St., Boston, MA 02115, USA

ARTICLE INFO

Article history:

Accepted 13 November 2013

Available online 22 November 2013

Keywords:

Enhanced permeation and retention effect

Active targeting

Nanoparticles

Nanomedicine

Personalized medicine

Tumor microenvironment

Drug delivery

Patient enrichment

Vessel normalization

Imaging

ABSTRACT

Cancer nanotherapeutics are progressing at a steady rate; research and development in the field has experienced an exponential growth since early 2000's. The path to the commercialization of oncology drugs is long and carries significant risk; however, there is considerable excitement that nanoparticle technologies may contribute to the success of cancer drug development. The pace at which pharmaceutical companies have formed partnerships to use proprietary nanoparticle technologies has considerably accelerated. It is now recognized that by enhancing the efficacy and/or tolerability of new drug candidates, nanotechnology can meaningfully contribute to create differentiated products and improve clinical outcome. This review describes the lessons learned since the commercialization of the first-generation nanomedicines including DOXIL® and Abraxane®. It explores our current understanding of targeted and non-targeted nanoparticles that are under various stages of development, including BIND-014 and MM-398. It highlights the opportunities and challenges faced by nanomedicines in contemporary oncology, where personalized medicine is increasingly the mainstay of cancer therapy. We revisit the fundamental concepts of enhanced permeability and retention effect (EPR) and explore the mechanisms proposed to enhance preferential "retention" in the tumor, whether using active targeting of nanoparticles, binding of drugs to their tumoral targets or the presence of tumor associated macrophages. The overall objective of this review is to enhance our understanding in the design and development of therapeutic nanoparticles for treatment of cancers.

© 2013 Elsevier B.V. All rights reserved.

Contents

1.	Introduction	3
2.	Passive targeting: nearly 30 years of the EPR effect	4
2.1.	The fundamentals of EPR	4
2.2.	Factors affecting the EPR effect	5
2.2.1.	Extravasation	5
2.2.2.	Diffusion and convection in the interstitium	6
2.3.	Tumor biology	6
2.3.1.	Tumor vasculature	6
2.3.2.	Tumor extravascular environment	6
2.3.3.	Improving EPR by changing tumor biology	7
2.4.	The physicochemical parameters	7
2.4.1.	Size	8
2.4.2.	Charge	8
2.4.3.	Shape	9
2.5.	The EPR effect in humans	9
2.6.	Future perspective on passive targeting	10
3.	Active targeting: toward a magic bullet?	10
3.1.	The fundamentals of active targeting	10

[☆] This review is part of the Advanced Drug Delivery Reviews theme issue on "Cancer Nanotechnology".

* Corresponding author at: Brigham and Women's Hospital, 75 Francis St., Boston, MA 02115, USA.

E-mail address: ofarokhzad@zeus.bwh.harvard.edu (O.C. Farokhzad).

¹ Both authors have contributed equally.

3.2.	Ligand conjugation/attachment strategies	11
3.2.1.	Pre-conjugation vs. post-formulation strategies	11
3.2.2.	Synthetic strategies for conjugation	11
3.2.3.	Non-covalent approaches	12
3.3.	Influence of the architecture of actively-targeted NPs	12
3.3.1.	The ligand density	12
3.3.2.	The NP size and shape	13
3.3.3.	Surface and ligand charge	13
3.3.4.	Surface hydrophobicity	13
3.4.	Targeting ligands	14
3.4.1.	Antibodies and their fragments	14
3.4.2.	Other proteins	14
3.4.3.	Peptides	15
3.4.4.	Nucleic acid based ligands	16
3.4.5.	Small molecules	16
3.5.	Active targeting in humans	17
3.5.1.	The choice of the target	18
3.5.2.	Assessing the impact of active targeting	18
4.	The near future of cancer nanomedicines	19
	Acknowledgments	19
	References	19

1. Introduction

More than 40 years ago, the foundations were laid down for nanotechnologies to deliver therapeutic and diagnostic agents in a safer and more efficient manner [1]. Achieving this vision became more realistic in recent years, with increasing numbers of nanotherapeutics and nanodiagnosics being commercialized or having reached clinical stage. In addition other important bench-to-bedside milestones are being achieved. In 2010, the first clinical evidence of gene silencing was obtained by systemically-administered targeted nanoparticles (NPs) delivering siRNA therapeutics [2]. Other clinical evidences of RNA interference have been obtained since then [3]. In parallel, clinical investigation of the first actively-targeted polymeric NPs, BIND-014, for the delivery of a small molecule drug (docetaxel) was reported [4]. Although only a relatively small number of nanosized drug delivery carriers have been approved for human use so far, it is now accepted that nanotechnologies will likely constitute a growing share of the oncologist's therapeutic arsenal over the next decades to come [5–7]. There are many nanoparticle technologies under development and a great majority are still without preclinical proof of concept. However, what is exciting is the momentum in this field: of the 81,000 articles on “nanoparticle” reported in Pubmed as of November 2013, more than half were published since 2010, emphasizing that research efforts are growing exponentially.

Nanoparticles offer the possibility to encapsulate poorly soluble drugs [8,9], protect therapeutic molecules [10], and modify their blood circulation and tissue distribution [11,12]. These properties are attractive in oncology in order to encapsulate cytotoxics exhibiting wide-ranging toxicities and physicochemical properties. For instance, liposome-encapsulated doxorubicin (DOX) decreases cardiac toxicity of the cytotoxic drug [13,14], and albumin-stabilized paclitaxel (nab-PTX) allows higher tolerated doses in patients [15]. Lately, drugs that modulate cancer signaling pathways (*i.e.*, molecularly targeted therapies) have shifted the paradigm of cancer treatment in patients exhibiting specific genetic mutations [16]. Like their cytotoxic counterparts these targeted drugs have toxicities and suboptimal tumor distributions that motivate their encapsulation in therapeutic NPs [17]. Furthermore, the robustness and redundancy of the signaling networks as well as the cross-talk between molecular pathways often promote resistance in cancers treated with molecularly targeted therapies [18,19]. For many kinase inhibitors on-target activity requires sufficient plasma concentrations of the drug and molecular pathways can reactivate as plasma concentrations decline [20]. Using NPs to precisely control the tumor levels of protein kinase inhibitors could theoretically circumvent that problem and result in improved efficacy.

Nanotechnologies are also appealing because they can facilitate the combination regimens which are commonly practiced in cancer therapy. Having a single NP encapsulating multiple active pharmaceutical ingredients (API) could potentially offer synergistic effects to promote the efficacy of therapies, while limiting the risk of resistance. When multiple drugs are administered separately, each API acts according to its own distinct pharmacology. Because drugs differ in their pharmacokinetic and pharmacodynamic properties, there is no certitude that target cells or tissues will synchronously receive optimal levels of each therapeutic entity. Conversely, when drugs are combined in a single NP carrier, the spatiotemporal exposure of each drug can be controlled more precisely and this may translate to a synergistic action among the API. The timely co-delivery to cancer cells of multiple agents inhibiting distinct, essential pathways could provide improved anticancer effects. This synergy has been demonstrated for combinations of small molecular weight drugs *in vitro* [21], combinations of a cisplatin prodrug and siRNA *in vivo* [22] or the combination of siRNA targeting 2 different genes in humans [3], highlighting the potential of encapsulating multiple API in a single carrier. Nevertheless, the determination of optimal therapeutic combinations using NPs is challenging. In opposition to conventional anticancer regimens where the dose of each single drug can be adjusted individually in patients (*i.e.*, based on their response or susceptibility to toxicities), the ratios of the different APIs encapsulated in a NP need to be optimized *a priori*, during the development phase. The large diversity of regimens possible makes the selection of ideal combinations difficult. Furthermore, temporal exposure to the different therapeutic components might also be important to maximize efficacy and avoid synergetic toxicities. Anticancer treatments undeniably impact transcriptional response in the tumor microenvironment and cancer cells, sometimes with important impact on subsequent response to chemotherapy (for example through p53 gene down-regulation or WNT16B expression) [23,24]. Combination regimens must therefore be designed so that the chronology of exposure to one agent does not affect the efficacy of the second drug. This might be particularly true for antiangiogenic agents where the shutdown of oxygen supplies has been shown to significantly reduce susceptibility of cancer cells to other chemotherapeutics [23]. Although it is feasible to independently control the release of APIs from a NP, the sequence of exposure must be specifically considered to maximize therapeutic synergism [25]. Robust chemistry, manufacturing and control processes also need to be devised to address the complexities of the multi-API therapeutics for successful development and commercialization.

Currently, despite the broad interest surrounding nanomedicines, the development and clinical translation of NPs remain laborious.

Among the clinically-validated nanomedicines, only nab-PTX has become officially part of the first-line treatment of cancer, but the picture is changing rapidly. Very recently, a combination of gemcitabine with nab-PTX proved to significantly improve survival from 6.7 months to 8.5 months (25% increase) in pancreatic cancer patients, as compared to gemcitabine alone [26]. These findings led, during the writing of this article, to the approval of nab-PTX as first-line treatment of advanced pancreatic cancer (in combination with gemcitabine). In all likelihood, the true potential of nanomedicines to really improve clinical outcome will continue to emerge as we begin to understand which patients can benefit further from treatment with anticancer therapeutic NPs. In the era of personalized medicine and as our understanding of genomics, tumor biology and nanotechnology progresses, sophisticated strategies may allow to address interpatient variability when initiating a treatment, and ultimately improve therapeutic response and cancer survival.

Molecularly targeted therapies have paved the way for the personalized treatment of patients based on the genetic profile of their tumors. It is now acknowledged that all patients do not respond equally to therapies [16]. Given that inter- and intratumoral variability can affect the architecture of the neovasculature and the tumor microenvironment, it becomes apparent that the passive targeting of NPs to tumors may be more complex than originally assumed [27]. The development of future therapeutics based solely on passive targeting strategies might not capitalize on the full potential benefit of therapeutic NPs. More than ever, increasing the response rates by screening patients for optimal response to nanomedicines based on specific tumor characteristics or the presence of certain biomarkers seems attractive, if not yet fully achievable. While patients who are naturally more responsive to NPs might receive more efficacious treatments, regimens increasing the susceptibility of the tumor to the nanomedicine could also be envisioned for others [28].

The parameters governing effective passive transport and retention of NPs in tumors hold true for actively-targeted NPs which are decorated with ligands. Although they have been proposed for quite some time [29–32], the first actively-targeted NPs have relatively recently made their way to human clinical trials [33]. This strategy exploits the affinity between surface ligands and antigens on the surface of cancer cells to facilitate the delivery of the NPs; actively-targeted NPs therefore display an increased degree of complexity. To potentially benefit from the active targeting strategy, it is imperative that the specific antigen be present and accessible on the targeted cells to bind the NPs. It is also important that antigen localization and expression remains adequate throughout the treatment. In this context, identification of predisposed patients goes beyond relatively simple genetic profiling.

Understanding the biological processes involved in the distribution and retention of nanomaterials inside the tumors is therefore essential to the development of personalized nanomedicine approaches. The concepts apply to therapeutic nanomaterials in general, whether drug carriers or therapy mediators exerting therapeutic effects in response to external stimuli (e.g., light [34], magnetic field [35,36], ultrasounds [37,38]). In this review, we examine the fundamentals behind targeting of nanomedicines to tumors and cancer cells. We will discuss how the morphology of the tumor vasculature, the components of the extracellular matrix or the presence of immune cells in the tumor microenvironments can affect the distribution of NPs. With the perspective of developing new therapeutic NPs, we will also examine how the physicochemical parameters of the nanomaterials affect their localization, retention, cell binding, internalization, efficacy and toxicity.

2. Passive targeting: nearly 30 years of the EPR effect

The observation that certain macromolecules accumulate preferentially to tumors was witnessed more than 30 years ago [39]. The tumor accumulation of therapeutic macromolecules was first

reported for poly(Styrene-co-Maleic Acid)-NeoCarzinoStatin (SMANCS), a 16 kDa polymer conjugate that non-covalently binds albumin in the circulation to reach a molecular weight of around 80 kDa [40,41]. The distribution of SMANCS to the tumor vicinity was observed in early preclinical development and led Matsumura and Maeda to further investigate the phenomenon [39]. Using labeled albumin and other proteins in addition to the polymer conjugate, they showed that proteins larger than 30 kDa (i.e., SMANCS, murine and bovine albumins (67 and 69 kDa, respectively) and IgGs (160 kDa)) could preferentially distribute to the tumor interstitium and remain there for prolonged periods of time [39]. This preferential distribution to the tumors was ascribed to the presence of fenestrations in the imperfect tumor blood vessels and to the poor lymphatic drainage in the tissue. The combination of these two phenomena was coined as the enhanced permeation and retention effect. Since then, the EPR effect has become the *leitmotiv* of many scientists for the efficient delivery of anticancer drugs to tumors, whether using polymer conjugates, liposomes or NPs.

Nonetheless, the EPR effect is much more complex than initially defined. The phenomenon has somehow become a blanket term encompassing dozens of complex biological processes (e.g., angiogenesis, vascular permeability, hemodynamic regulation, heterogeneities in tumor genetic profile, heterogeneities in the tumor microenvironment and lymphangiogenesis). These factors vary among patients and their tumor types. Likewise, the distribution and accumulation of NPs in tumors are also multifaceted and are affected by the biological and physicochemical properties of each material. For these reasons, and with the advent of personalized medicine, the designation of EPR as a simple self-explaining phenomenon might be becoming outdated.

2.1. The fundamentals of EPR

When a solid tumor reaches a given size, the normal vasculature present in its vicinity is not sufficient to provide all the oxygen supply required for its further proliferation. As cells start to die, they secrete growth factors that trigger the budding of new blood vessels from the surrounding capillaries [42]. This process, known as angiogenesis, promotes the rapid development of new, irregular blood vessels that present a discontinuous epithelium and lack the basal membrane of normal vascular structures [43,44]. The resulting fenestrations in the capillaries can reach sizes ranging from 200 to 2000 nm, depending on the tumor type, its environment and its localization [45]. When blood components reach the abnormal, discontinuous vascular bed, the fenestrations offer little resistance to extravasation to the tumor interstitium. This denotes the enhanced permeation portion of the EPR effect.

In normal tissues, the extracellular fluid is constantly drained to the lymphatic vessels at a mean flow velocity around 0.1–2 $\mu\text{m/s}$ [46]. This allows the continuous draining and renewal of interstitial fluid and the recycling of extravasated solutes and colloids back to the circulation. In tumors, the lymphatic function is defective, resulting in minimal uptake of the interstitial fluid [47]. As a result, the colloids cannot rely on convective forces to return to circulation. While molecules smaller than 4 nm can diffuse back to the blood circulation and be reabsorbed [48–50], the diffusion of macromolecules or NPs is hindered by their larger hydrodynamic radii. Therefore, NPs that have reached the perivascular space are not cleared efficiently and accumulate in the tumor interstitium. This aspect represents the enhanced retention component of the EPR effect.

Since the early works of Matsumura and Maeda in the mid-1980s, the EPR effect has been comprehensively documented using various tumor types and animal models. The parameters which affect the distribution of macromolecules and NPs to the tumor are better understood, and we are slowly unraveling the subtleties of the EPR effect [44,51]. Importantly, it is now recognized that lymphatic drainage is not homogeneous throughout the cancerous mass. Vessels in the bulk of the tumor experience higher mechanical stress, and the functional loss in the

intratumoral regions is therefore more important than in the margin [47]. In fact, residual lymphatic activity and *de novo* lymphangiogenesis are believed to be in part responsible for the progress and dissemination of metastases [52]. The heterogeneity of lymphatic function within the tumor is therefore a factor that should be considered when addressing tumor NP accumulation.

2.2. Factors affecting the EPR effect

In a solid tumor, the distribution of molecules to the tumor is governed by at least three distinctive but interrelated phenomena: the extravasation of colloids from the blood vessels, their further diffusion through the extravascular tissue and their interaction with intracellular and/or extracellular targets within the tumor environment (Fig. 1). The first two aspects are the result of diffusive and convective forces and can be influenced concurrently by the tumor biology and the characteristics of the diffusing species. The third parameter is more complex and less understood. It represents the interactions of the colloids with the tumor whether through adsorption phenomena, cellular uptake or degradation and metabolism. These aspects can all affect the equilibrium of accumulation inside the tumor; they depend on the nature of the material, its affinity for all components of the tissue and the tumor composition (e.g., nature of the extracellular matrix, the type of cells present). Because we still lack the full understanding to completely assess all these interactions, they are represented as a “black box” that will be further detailed below (Table 1).

2.2.1. Extravasation

Extravasation of colloids from the circulation is influenced by their concentration in the blood, the relative permeability of the vascular wall to macromolecules and NPs as well as the nature of the extravascular environment. In order to reflect the relative involvement of each factor on the escape of colloids from the blood, the phenomenon can be described by simple and general mathematical relations. Although these equations are difficult to use experimentally because of the

Table 1

Extravasation as well as interstitial diffusion and convection can be affected by the tumor biology and the various characteristics of the colloid.

Tumor biological properties	Parameters potentially affected
Vessel architecture (e.g., fenestrations, blood flow)	P, A, P _v , L _p , σ, σ _F Black box (endothelial uptake)
Interstitial fluid composition	π _i , Black box (protein adsorption)
Extracellular matrix composition	D _{eff} , R _i , π _i , Black box (adsorption)
Phagocyte infiltration (e.g., TAM, dendritic cells)	R _i , Black box (cellular uptake)
Presence of necrotic domains	P _i , π _i , D _{eff} , R _i Black box (protein adsorption, colloidal stability)
<i>NP properties</i>	
Blood circulation times	C _v , C _i
Size	P, L _p , D _{eff} , σ _F , φ _i , R _i Black box (colloidal stability, cellular uptake)
Charge	P, L _p , D _{eff} , R _i Black box (colloidal stability, cellular uptake)
Shape	P, L _p , D _{eff} , σ _F , φ _i , R _i Black box (adsorption, cellular uptake)
Surface characteristics (e.g., hydrophobicity, ligands)	P, L _p , R _i Black box (adsorption, cellular uptake)

great number of variables involved, they can be useful to represent and understand the consequences of each biological and physicochemical characteristic independently.

The total flux of material toward the tumor is described by adding the contributions of diffusive (in red) and convective forces (in blue) as well as possibly a number of unknown phenomena (black box):

$$J_{Total} = PA(C_v - C_i) + L_p A [(P_v - P_i) - \sigma(\pi_v - \pi_i)](1 - \sigma_F) C_v + \text{BLACK BOX.} \tag{1}$$

The diffusive component (PA(C_v - C_i)) originates from the Brownian motion of colloids and results in a positive net flux toward the interstitium when a gradient exists between the vascular (C_v) and interstitial concentrations (C_i) [46]. It is given by a modification of Fick's first law to account for the permeability of the vascular wall (P) and the area of the blood vessel (A). The permeability incorporates both the diffusion coefficient of the colloid (D) and how the vascular barrier restricts its passage. This hindrance effect depends on the physicochemical properties of the colloid as much as on the properties of the vessel wall [44].

The outflow of fluids from the vessel also creates a convective force pushing the colloids toward the tumor. In that case the flux of fluid (L_pA[(P_v - P_i) - σ(π_v - π_i)]) is described by the Starling Law where L_p is the filtration coefficient of the fluid through the vessel wall, P_v and P_i are the vascular and interstitial hydrostatic pressures, π_v and π_i are the vascular and interstitial oncotic pressures and σ is the capillary osmotic reflexion coefficient [46]. This later parameter reflects the permeability of the capillary to proteins and how effective the oncotic pressure gradient is to pull the fluid back in the vascular space. The flux of fluid is adjusted for the imperfect permeation and drag of the colloid by the fluid (σ_F) and the colloid concentration in the vascular compartment (C_v).

Finally, despite the recent progresses in cancer biology and nanotechnology, our understanding of the tumor microenvironment and how colloids extravasate and reach the tumor cells is far from complete. The “black box” in Eq. (1) highlights this opportunity for further exploration. When measuring the tumor uptake of a compound, the distinction between extravasated material in the vicinity of the vessels and




Fig. 1. The EPR effect results from 2 distinct phenomena: the extravasation of the colloid from the blood vessels and their subsequent movement in the tumor extracellular matrix (ECM) by diffusion and convection.

the material which is taken up by tumor endothelial cells in the neovasculature is difficult. Some researchers describe enhanced interactions with endothelial cells, for example by cationic charges (see Section 2.4.2), as an increase in the effective P of the vessel [53,54]; others consider these interactions to be the product of absorption and endocytosis by the endothelium [55,56]. The black box accounts, among other things, for that uncertainty on the effective concentration in the vasculature available for extravasation (e.g., if endothelial cells are internalizing the colloids relatively fast, the actual concentration left to permeate through the vessel is lower than the concentration supplied by the systemic circulation, C_v).

Similarly, authors have shown that tumor accumulation of NPs increased with higher levels of phagocytic cells in the tumor interstitium [57]. The interactions of NPs with macrophages and dendritic cells can therefore affect the concentrations of NPs effectively diffusing in the tumor microenvironment (C_i). Finally, it is also important to decouple the accumulation of the NPs by EPR effect from the accumulation of their encapsulated payload. The tumor distribution of therapeutic NPs is usually measured by tracking their loaded cargo, whether it is a small molecular weight drug, a therapeutic macromolecule, a fluorescent dye or a radiolabeled tracer. Those entities may have very distinct properties when they are encapsulated in NPs compared to when they are released; the kinetics of their release as well as their possible interactions with the tumor micro-environment must therefore be accounted for.

The black box is introduced to address all parameters which factor in the retention of the NPs in the tumor whether they are the product of absorptive or intracellular trafficking forces or the result of degradation, metabolism and drug release. It also acknowledges that many aspects of tumor accumulation are still not fully understood.

2.2.2. Diffusion and convection in the interstitium

Once the colloids are extravasated to the tumor, their surrounding environment is composed of interstitial fluid (with a composition similar to plasma), cancer and stromal cells and the extracellular matrix (ECM). Their movements in the interstitium are also guided by convective, diffusive forces and other phenomena and can be described by:

$$\frac{\partial C_i}{\partial t} = D_{\text{eff}} \nabla^2 C_i + \varphi_i \underline{v} \cdot \nabla C_i - R_i \quad (2)$$

Here, the changes in interstitial concentration over time ($\delta C_i / \delta t$) result from the diffusive component ($D_{\text{eff}} \nabla^2 C_i$), the convective component ($\varphi_i \underline{v} \cdot \nabla C_i$) and the effects of the tumor microenvironment on the colloid's transport (R_i). The diffusive factor is governed by the effective diffusion coefficient (D_{eff}) and the change in concentration in all directions (i.e., the Laplace operator of the interstitial concentration, $\nabla^2 C_i$). The convective portion results from the direction and intensity of the convective motion (i.e., the fluid velocity vector (\underline{v})), the spatial concentration gradient (∇C_i) and a coefficient (φ_i) to account for the fact that the colloid velocity in the porous ECM might be different from the velocity of the fluid (e.g., drag, adsorption or exclusion effects). This difference can be explained by interactions with the ECM retaining the colloid ($\varphi_i < 1$) or by size-exclusion effects hastening it through larger pores ($\varphi_i > 1$). Finally, the factor R_i also highlights the possible degradation, metabolism or binding of the colloids to ECM, their capture by extravasated components of the mononuclear phagocyte system (MPS) as well as their uptake by tumor cells. The negative sign acknowledges that higher interactions with the tumor result in decreased movement.

2.3. Tumor biology

The tumor environment is a complex milieu where normal principles of anatomy and physiology are defied. The untamed growth of cancer cells and the secretion of angiogenic factors both contribute to the highly disorganized vasculature and congested extravascular

environments. Paradoxically, these structural imperfections are both the cause of the EPR effect and the principal source of uneven tumor accumulation and retention of nanomaterials.

2.3.1. Tumor vasculature

The release of angiogenic factors from cancer cells dictates the blood vessel architecture and morphology of the vascular wall [42]. The neovasculature sprouting from capillaries is disordered, discontinuous and highly fenestrated [28]. The degree of leakiness of the endothelium and the enhanced vascular permeability to macromolecules and nanomaterials depend on many factors like the cancer type, its stage and in xenograft models, the implantation site of the tumor [44,45,58]. In Eq. (1), the vascular biology affects both the diffusion (via the permeability, P) and the convection through the vascular wall (via the filtration coefficient (L_p)).

While the fenestrations offer escape routes for the colloids, the discontinuities and irregularities in the architecture of the vessels also affect the blood flow and the hydrostatic pressure in the vessels [28,44]. The mass of hyperproliferative cancer cells also exert mechanical pressure on the different vessels further impeding perfusion [47]. The reduced pressure might be associated with a decrease in the convective forces responsible for the extravasation of blood components and nanomaterial ($\downarrow P_v$).

2.3.2. Tumor extravascular environment

The tumor extravascular environment is a congested entanglement of collagen fibers and glycosaminoglycans (GAG), with inhomogeneous distributions of solutes, proteins and cellular debris. In opposition to healthy tissues where interstitial flow is regulated to efficiently disseminate the fluid throughout the cell population [46,49], the disorganized structures in tumors strongly hinder the diffusion and fluid convection. The abnormal traffic of fluid influences local interstitial hydrodynamic (P_i) and oncotic pressures (π_i), two parameters which impact the convection of NPs through the vascular wall [59].

Once NPs have extravasated, the nature of the ECM further regulates the diffusive and convective movement of NPs in the tumor. Various studies *in vivo* and *ex vivo* have shown that the effective diffusion coefficient (D_{eff}) in the tumor interstitium is below that measured for colloids in simple solutions [60,61]. The GAGs (e.g., hyaluronic acid, heparan sulfate), covalently-attached to proteins (e.g., collagen, laminin), affect the viscosity of the environment and the tortuosity of the diffusive paths. GAG chains organized in blotches of low and high viscosities which translate into a two-phase transfer process where colloids of different sizes show high and low mobilities, respectively (Fig. 2, closed and open symbols) [61]. Although these distinct mobilities can probably be explained by differences in the diffusion of colloids (D_{eff} , in Eq. (2)), the presence of some steric exclusion effects (φ_i) caused by the inhomogeneous environment cannot be completely ruled out.

The collagen content and its degree of organization in the ECM also correlates with the resistance exerted on the interstitial transport ($\downarrow D_{\text{eff}}$) [62]. The disruption of the collagen network by collagenase can break up the protein entanglement and restore some mobility to slow diffusing species [61,62]. Similarly, intratumoral injections of collagenase were shown to rearrange the ECM in a manner that allowed enhanced mobility of 150-nm viral vectors in the tumor [63], further consolidating the influence of diffusion on the two-phase transport (in opposition to steric effects).

On the other hand, the effects of GAG-disrupting enzymes are less clear. In xenograft models of various tumors, relations between the GAG content and the transport restriction could not be established [62]. In some instances, hyaluronidases were reported to decrease the diffusion of macromolecules by collapsing the hydrated protein structure and increasing the overall viscosity ($\downarrow D_{\text{eff}}$ and φ_i) [61]. In other cases, heparinases that cleave the matrix heparan sulfate moieties can restore mobility of positively-charged macromolecules [60], in this




Fig. 2. After intratumoral injection in melanoma xenografts (Mu89), the diffusion coefficient of macromolecules and nanomaterial in the ECM is inversely proportional to the hydrodynamic radius. Diffusion of bovine serum albumin (BSA, circles), 70-kDa dextran (squares), immunoglobulins (IgG, triangles), 2-MDa dextran (diamonds) and liposomes (inverted triangles) follows 2 phases with fast (closed symbols) and slow colloid populations (open symbols). Adapted with permission from [61].

case most probably by decreasing the adsorptive interactions of the colloids with the ECM ($\downarrow R_i$ in Eq. (2)).

Finally, cells from the MPS have been found to extravasate to the tumor interstitium. The affinity of these macrophages for the colloid and the resulting phagocytic activity can further impede the NPs' movement toward the cancer cells while increasing their retention in the tumor interstitium [27]. Recently, Zamboni and colleagues showed that xenografts of ovarian cancer showing increased amount of CD11c-positive cells (a marker specific for dendritic cells) had increased liposomal accumulation compared to melanoma cells with lower dendritic cell expression [57]. Although it is unclear if the increased tumor accumulation seen in this study has a positive or negative influence on the NPs' therapeutic efficacy, it certainly implies that the MPS plays a role in their retention in tumors (decreasing movement by $\uparrow R_i$, in Eq. (2)). In humans, the age of patients and their intrinsic MPS activity were also correlated with the clearance of liposomes from the blood [64]; older patients or patients with hepatic metastases were shown to experience higher blood exposure to the NPs. Interestingly, older patients also had less hematologic toxicity compared to patients below 60 years old [65], further suggesting that interactions between NPs and the MPS affect the pharmacodynamics of nanomedicines. Nevertheless, this phase 1 clinical trial did not focus on how the different MPS functions in the tumor correlate with the efficacy of the NPs. To answer this question, a new clinical trial was recently initiated to use iron oxide NPs as an MRI contrast agent in combination with therapeutic liposomes (clinicaltrials.gov, NCT01770353). This pilot study will be discussed in more detail in Section 2.5.

2.3.3. Improving EPR by changing tumor biology

As our knowledge about the EPR effect grows, pharmacological approaches are proposed to optimize the tumor microenvironment for enhanced distribution of macromolecules and nanomaterials. The injection of enzymes that remodel the ECM to augment the intratumoral mobility of colloids ($\uparrow D_{\text{eff}}$) has been introduced before [63]. In a similar fashion, other methods remodeling the perivascular environment have been exploited using photo-immunotherapy [66] or small molecular weight drugs [67]. These approaches showed increased tumor

distribution and therapeutic efficacy of oncolytic viruses [63,67] and/or NPs [66,67] in preclinical models.

Another approach relies on improving the transvascular convective movement by increasing the perfusing pressure (P_v). To that end, the cyclic or continuous administration of hypertensive molecules, like angiotensin II, was studied [59]. This approach resulted into significantly increased extravasation only when colloids (in this case, antibodies) had sufficient affinity ($\uparrow R_i$) to bind to the tumor and avoid being translocated back to the circulation upon pressure normalization. The administration of enalapril, an angiotensin-converting enzyme (ACE) inhibitor, was also proposed to augment tumor accumulation of antibodies [68]. In this case, the ACE inhibitor was used to block the degradation of bradikinin, a potent physiological vasodilatory peptide that increases the vascular permeability to macromolecules ($\uparrow P$ and L_p). The combination of enalapril with angiotensin II was effective to counter the hypotensive effect of the ACE inhibitor and further improve EPR. Other vasodilatory pathways ($\uparrow P$ and L_p), like nitric oxide [69], prostaglandins [70] or carbon monoxide [71] were also found to improve tumor accumulation of NPs.

The normalization of the blood vasculature was also proposed as an alternative to improve the EPR effect [28]. This approach is believed to rectify the blood flow in the tumor ($\uparrow P_v$) and consequently normalize the interstitial fluid exchanges ($\downarrow P_i$ and π_i). In a murine model of orthotopic allograft breast adenocarcinoma, the blocking of the vascular endothelial growth factor receptor-2 (VEGF-R2) improved penetration of small (~ 12 -nm) albumin-based drug complexes while not affecting the diffusion of larger (> 60 nm) nanoparticles [72]. It is believed that the restoration of the transvascular pressure gradient also impacted the size of the endothelial fenestrations and resulted in decreased permeability for the larger particles ($\downarrow P$ and L_p).

The impact of the modulation of the vessel morphology strongly depends on the initial tumor architecture. The administration of various small molecular weight tyrosine kinase inhibitors affecting the vascular endothelial growth factor (VEGF) activation pathways had different effects in murine xenograft models of colon and pancreatic cancers [73]. In the former, VEGF-inhibition lead to decreased permeability by reducing the endothelial density in the tumor ($\downarrow A$ and possibly P and L_p), while in the latter, it decreased the coverage of the vessels by pericyte and improved permeability ($\uparrow P$ and L_p). The administration of transforming growth factor- β inhibitors (TGF- β inhibitors) in a model of hypovascular, poorly-permeable pancreatic cancer, was also found to increase the permeability of blood vessels to large nanoparticles ($\uparrow P$ and L_p) [73–75].

Together, these results in preclinical models highlight certain therapeutic interventions that could possibly modulate the tumor biology and regulate the EPR effect. However, further efforts are needed to fully understand how to efficiently exploit these different strategies in a clinical setting.

2.4. The physicochemical parameters

The physicochemical parameters of the colloids affect their extravasation by influencing, among other things, their diffusivity (D_{eff}), their permeability through the vascular wall (P and L_p), their drag coefficient in the fluid (ρ_f and φ_i) and their interactions in the tumor with the cells and the ECM (R_i). Furthermore, the physicochemical characteristics of exogenous materials used for therapeutic or diagnostic applications also impact on how the host's defense mechanisms clear them from the blood circulation [11,12]. In fact, because individual physicochemical parameters can affect the overall blood circulation kinetics, the extravasation processes and the intratumoral diffusion, directly measuring the influence of each specific characteristic on the EPR is difficult.

Nevertheless, the total blood exposure to the NP is believed to be a key factor influencing its distribution to the tumor in the EPR effect [76]. The diffusive and convective elements forces are both influenced

by the concentration of colloid in the bloodstream (C_v). Maintaining high blood concentrations is also very important to ensure unidirectional diffusion toward the tumor and prevent the efflux from the tumor when $C_i > C_v$ [77]. Therefore, longer circulation times in the blood result in higher amounts extravasated to the tumor interstitium. The tumor deposition of soluble polymers like poly(ethylene glycol) (PEG) and N-(2-hydroxypropyl)methacrylamide (HPMA) augments proportionally to blood exposure when the molecular weight is increased above the glomerular filtration threshold and the polymer cannot be eliminated by the kidney (Fig. 3) [50,78]. The correlation between the tumor accumulation and the blood circulation kinetics has since then been generalized to other polymers [79], liposomes [80,81] and nanoparticles [82].

2.4.1. Size

The size of the nanomaterial influences the kinetics and extent of tumor accumulation. The material needs to be smaller than the cut-off of the fenestrations in the neovasculature, but size is also factored in the various parameters affecting extravasation to the tumor and diffusion in the interstitium (P , L_p , D_{eff} and, possibly to a lesser extent, ρ_F and φ_i). In mice xenograft models, when the kinetics of intratumoral accumulation were studied over 30 min, smaller macromolecules (3.3 and 10 kDa, R_H : 2 and 3 nm) were shown to accumulate faster and diffuse deeper in the tumor than larger molecules (70 and 2,000 kDa, R_H : 7 and 25 nm) [77]. However, this accumulation was transitory as smaller molecules would rapidly diffuse back in the vascular compartment [77]. By focusing on a short accumulation period during which the blood concentrations of each polymer were relatively constant, the authors of this study could somehow alleviate the influence of the distinct circulation times on their findings.

A recent article addressed this issue by comparing the tumor deposition of different-sized particles with similar circulation profiles [74]. This study found that particles (sizes: 30, 50, 70 and 100 nm) distributed comparably when the tumors were hyperpermeable (murine colon adenocarcinoma), but that only NPs smaller than 70 nm could accumulate efficiently in poorly permeable tumors (human pancreatic adenocarcinoma). In the latter case, the degree of tumor accumulation of these platinum-loaded particles correlated with tumor shrinking efficiency. This study highlights the dual effect of the tumor biology and the size of nanomaterial on the endothelial permeability (P and L_p) and nanomaterial reflexion coefficient (ρ_F). Given the relatively small difference in the hydrodynamic radii of the NPs, it suggests that the neovasculature wall in this pancreatic cancer model acts as an all-or-nothing barrier with a 30 to 70 nm threshold (P and $L_p = 0$, when




Fig. 3. In murine S-180 sarcoma, the tumor accumulation levels of macromolecules (○) are in direct relation with the total body exposure (AUC, ▲) and inversely proportional to their renal clearance (●). This holds true for other types of tumors and nanomaterials. Used with permission from [76].

size > threshold). This cut-off could be raised above 70 nm when the tumors were treated with TGF- β inhibitor (see Section 2.4.3). Preferential tumor accumulation of smaller particles (*i.e.*, <50 nm) was confirmed by others, however differences in the blood circulation profiles of the compared NPs could not rule out other interfering parameters [83].

Insightful results were also obtained by Wong *et al.* who developed 90-nm NPs that disintegrated into 10-nm quantum dots when degraded by tumor proteinases [84]. The evaluation of their novel nanomaterial design emphasized the importance of size on the interstitial diffusivity of particles (D_{eff}). The quantum dots loaded in cleavable particles showed improved spatial distribution and enhanced depth of penetration in the tumor compared to quantum dots loaded in similarly-sized, non-cleavable silica NPs.

2.4.2. Charge

Like other physicochemical parameters, the charge of macromolecules [85] and nanomaterials [11,12] alters both systemic circulation times and intratumoral processes and addressing both phenomena independently is difficult [54]. The presence of surface charge can alter the opsonisation profile of the material, its recognition by cells in the organs of the MPS and its overall plasma circulation profile ($\downarrow C_v$) [11,12,86–92]. Negative surface charges can either increase, decrease or have no impact on the blood clearance of NPs [92–97], but positive charges are generally recognized as having a negative effect on the plasma exposure to the nanomaterial [90,91,98].

In tumor-bearing animals, despite the reduced blood circulation times, non-PEGylated, positively-charged liposomes containing the lipid 1,2-diacetyl-trimethylammonium propane (DOTAP) display higher ratios of concentration in tumor vs. surrounding tissue compared to their negative or neutral counterpart [55,56,98]. The preferential distribution to the tumor is attributed to localization of the vesicles to the epithelium of the tumor neovasculature, with very little extravasation or very shallow interstitial diffusion. The positive charges possibly favor interactions of the NPs with the tumor blood vessels and eliminate their predisposition to diffuse deeper in the tumor while preventing their redistribution in the systemic circulation. This phenomenon has been utilized for therapeutic purposes by targeting the tumor vessel endothelium with antiangiogenic drugs in preclinical models [55,99] and, very recently, in humans [100,101].

Sterically-stabilized colloids with positive charges have also demonstrated enhanced tumor accumulation [102,103] while others seem less efficacious [90,91]. Parameters like the degree of ionization, the relative blood circulation times of the control NPs [102] or the architecture of the construct (*i.e.*, charge on the core surface [102,103] vs. charge on the corona [90,91]) might explain these conflicting findings. Recent work has highlighted that the architecture of charge presentation by a zwitterionic material influences the type of non-specific interactions with endothelial cells [104].

Besides the limited extravasation of NPs due to interactions with the tumor endothelium, charges can also bind to the ECM and limit diffusion in the interstitium ($\uparrow R_i$) [60]. *Ex vivo* studies conducted on ECM isolated from mice sarcoma showed that charges (positive or negative) had deleterious effects on the movement of NPs through the matrix. In fact, the presence of charge above a certain threshold (>30% DOTAP or >60% 1,2-dioleoyl-*sn*-glycero-3-[phospho-*rac*-(1-glycerol)] (DOPG), *i.e.*, ζ -potential $\sim +10$ mV and -30 mV, respectively) abolished the diffusion of the NPs [60]. This is in accordance with the results obtained from the intratumoral injection of neutral and charged NPs, which showed that charged colloids interacted with the tumor longer than their neutral counterparts [105,106]. *Ex vivo*, the mobility of the particles can be restored by masking the surface charges with high amounts of salt (*e.g.*, adding 1 M KCl) or, in the case of the positively-charged material, small amounts of polyanionic proteins (*e.g.*, 0.5 mM Heparin), suggesting that this effect might be the result of ionic interactions between the NPs and the ECM [60].

2.4.3. Shape

The shape of NPs has been shown to influence blood exposure by modulating interactions with the MPS [12,107–109]. Furthermore, single-wall carbon nanotubes with high aspect ratio (*i.e.*, from 100:1 to 500:1) were shown to be cleared efficiently by the kidneys despite dimensions (*i.e.*, 100–500 nm) 10–20 times above the usual glomerular filtration threshold, suggesting that elongated shapes could provide benefits to the filtration process through porous structures [110]. These factors prompted investigation into the ability of nanomaterials with different aspect ratios to accumulate in the tumors.

Recently, the tumor distribution kinetics of nanorods with a length of 44 nm (aspect ratio: 10) were compared to those of 35-nm nanospheres showing the same hydrodynamic radius [111]. Despite similar blood circulation profiles, the nanorods were shown to extravasate to the interstitium 4 times faster and to diffuse deeper in the tumor, interacting with a 1.7-fold larger volume. Similarly, regardless of their increased splenic clearance and shorter blood circulation times, elongated viral nanofilaments derived from plant viruses showed increased tumor homing and accumulation compared to other spherical viral constructs [112]. All together, these findings suggest that elongated shapes might present beneficial EPR properties (possibly by influencing P_i , L_p and ϕ_i).

2.5. The EPR effect in humans

Although the EPR effect has been observed and studied in many rodent models using subcutaneous and orthotopic xenografts, as well as genetically-engineered mouse models, its prevalence in human primary and metastatic tumors requires further investigations [113,114]. This is particularly important since only a few first-generation anticancer nanomedicines materially improve overall survival in patient cohorts, conceivably because subpopulations with higher susceptibility to NPs might be masked by non-responsive patients [27]. Provided that such subpopulations exist, careful patient selection based on the degree of EPR could potentially further enhance the effectiveness of nanomedicines in clinical practice [27].

Many reasons can account for the paucity of data for the EPR effect in human, the most important being that obtaining meaningful biodistribution data in humans is challenging. In clinical studies, surrogates to the analytical methods employed in animals (*i.e.*, tissue extraction and digestion) must be employed [115]. During the initial evaluation of PEGylated liposomes as drug carriers for DOX and other drugs, some studies assessed the distribution of liposomes and their payload in patients' tumors [116–121]. These studies measured the drug concentration in malignant exudates [116], tumor biopsies [117,118] or using radionuclide-based imaging of modified liposomes (^{99m}Tc or ^{111}In) [119–121].

These studies usually report advantageous drug tumor deposition of the liposomes compared to the free DOX: the DOX levels achieved in the tumor 3 to 7 days after the administration are usually 4–16-fold higher with the liposomes [116,117]. This difference might be due to preferential accumulation of the liposomes in the tumor by EPR or to fact that liposomes maintain higher DOX concentrations in the blood, supplying the tumor for a longer time (free DOX $t_{1/2}$: 10.4 h, liposomal DOX $t_{1/2}$: 45.9 h).

In opposition to the animal studies where EPR of PEGylated liposomes can be evidenced by high tumor-to-blood ratios [122,123], Fig. 4 shows that, in humans, the only cancers which show preferential tumor accumulation for liposomes (*i.e.*, ratio > 1) are sarcomas (with the exception of 1 patient with bone metastases for which plasma and biopsy samples were not analyzed at the same laboratory) [118]. Evidently, the direct comparison of these studies is imperfect because the methodologies vary and the studies do not provide information on the antitumor response [64,124]. Also, it is still unknown how the data obtained with liposomes translates to other systems as factors like drug release from the NPs and cellular uptake might affect the drug




Fig. 4. In human, the accumulation of liposomes in tumors is different in each type of cancer. Sarcomas seem to be the sole cancers for which the accumulation of liposomes in tumor is superior to that in plasma or surrounding tissue. (●) From [116], DOX concentration in malignant infusion vs. plasma concentration, using Doxil®. (□) From [119], ^{111}In -DTPA-labeled liposomes in tumor ROI vs. plasma concentration. (◇) From [121], ^{99m}Tc -DTPA-labeled Doxil® tumor ROI vs. skull bone marrow. (▲) From [117] DOX concentration in tumor biopsies vs. plasma concentration, using DOX-containing PEGylated liposomes. (○) From [120], ^{99m}Tc -DTPA-labeled liposomes tumor ROI vs. surrounding tissue. (■) From [118], DOX concentration in bone metastases vs. plasma concentration, using Doxil®. All ratios are given for concentrations measured at the same time-point and individual patients are presented when possible; the doses of DOX used are labeled on the figure (closed symbols); open symbols represent the use of empty liposomes.

concentrations measured in the tumor. Nevertheless, these observations support the idea that diverse cancer types might show distinct predisposition to the EPR effect.

It is also important to note that systemic chemotherapy is usually reserved for patients with metastatic cancers, while preclinical models usually focus on primary tumors. Since primary and metastatic lesions may exhibit different behavior [113,114], it is still not clear if the deposition of NPs in metastases correlates to that of primary tumors [125]. To further complicate the picture, patients enrolled in early clinical phases are rarely naive to therapeutic agents, and most have received multiple treatments that may have altered the complex tumor microenvironment.

Finally, another reason why the assessment of the EPR effect in humans is not simple is that the optimal parameters to ensure maximum distribution and retention in humans can be different to those required in animal models. Both clearance mechanisms and tumor biology differ between animals and man [126,127]. Animal models, especially xenograft models, used to study NPs at the preclinical stage, seem to offer limited predictive value of the clinical outcome in humans [128]. Furthermore, while the screening of NPs with different physico-chemical properties in mice is relatively easy, the parallel evaluation of multiple NP candidates in humans remains largely impractical. Phase 0 clinical trials allow the concomitant study of several drug candidates (including NPs) at sub-therapeutic doses in a small number of patients to achieve a proof of principle [129]. However, this approach is potentially complex and would require the manufacturing of clinical supplies and suitable preclinical toxicology studies for each NP. Furthermore, assessment of tumor accumulation would also necessitate the development of robust endpoint assays, with adequate sensitivity regardless of the sub-therapeutic doses used and

sufficient reproducibility considering the low number of patients. Hence, while possible, the parallel evaluation of multiple formulations in humans remains a considerable challenge.

In the context of personalized medicine, it therefore becomes very important to identify what are the characteristics that predispose cancers to the accumulation of available NPs. In animals, iodine-loaded liposomes as EPR imaging modalities were able to determine *a priori* which mice would benefit from the injection of therapeutic nanomaterials [130]. A similar strategy in humans may provide useful information by correlating EPR effect with biomarkers and response to therapeutic NP, highlighting the potential role of clinical imaging in the development of nanomedicines. A pilot clinical study was recently initiated to ascertain the safety of the concomitant administration of an imaging agent with irinotecan-loaded liposomes (clinicaltrials.gov, NCT01770353). In this study, ferumoxytol, a clinically validated iron-oxide NP approved for the treatment of iron-deficiency, is being administered prior to the administration of MM-398, a liposomal irinotecan. Subsequently, intratumoral concentration of irinotecan and SN-38 (the active metabolite of irinotecan) will be measured. Given that ferumoxytol is efficiently phagocytosed and has been used to image tumor-associated macrophages (TAM) in preclinical models of solid tumors [131], this study might reveal insightful information about the possible predictive value of iron-oxide NP accumulation on the efficacy of MM-398. Similar iron-oxide NPs have been efficiently used in humans for the early detection of lymphatic metastases [132] and to measure increased vascular permeability in the context of inflammation [133].

2.6. Future perspective on passive targeting

The last 30 years have shown that EPR plays an important role in the delivery of macromolecules and nanomaterials to tumors. Nevertheless, a full understanding of the degree of interpatient variability and the importance of tumor heterogeneity in the EPR effect in humans has yet to be established [27]. As we unravel which parameters are crucial to improve the distribution of chemotherapeutics to tumor cells, the individual patients and the cancer types benefiting the most from carrier-mediated drug delivery might be identified.

Furthermore, it is now appreciated that the physicochemical properties of the nanomaterial are as important as the tumor biology. Meaningful exploitation of the EPR effect in humans will require precise understanding and control of the physicochemical properties of nanomaterials [134] and, possibly artificial fine-tuning of the tumor microenvironment [28]. In the era of modern cancer biology, assessing the tumor microenvironment in individual patients and predicting their susceptibility to the EPR effect may eventually become the mainstay of therapy when choosing between therapeutic regimens. Although it is hard to currently predict which markers better correlate with NPs efficacy in humans, relationships will most likely delineate as tumor genotyping and bioassays continue to progress. Ideally, a broad biomarker correlating the susceptibility of tumors to nanomedicines in general (*i.e.*, levels of tumor-associated macrophages; levels of ECM proteins/GAG; tumor infiltration of specific proteins/cells; endothelial expression of surface biomarkers) would certainly facilitate the screening of patients to optimize therapy.

In parallel, the EPR phenomenon affects the distribution of the drug-carrier to the tumor without necessarily increasing the ability of the drug to reach its pharmacological target. The optimization of NPs' efficacy therefore also involves optimal drug release rates whether through controlled diffusional release [135], covalent conjugation of the drug to a polymer backbone [136,137] or environment-triggered release [138,139]. Without specific affinity of the nanomaterial for the cancer cells, the chemotherapeutic payloads will have to reach their pharmacological targets by their own means or risk diffusing back into the vasculature [77]. In some instances, drugs exhibit sufficient affinity for their pharmacological target to remain trapped in the tumor for prolonged

periods of time. For example, the high affinity of docetaxel for the microtubules translates into very low efflux from the tumor; consequently, the elimination half-life of the drug from tumors is approximately 15–20 times higher than its elimination from the blood and normal tissue [140]. The behavior of drugs and their affinity for the intratumoral environment need to be taken into account individually when designing passively-targeted NPs and the optimal drug release profiles should be optimized on a case-by-case basis. In this regard, tools that can assess the interactions between drugs and their substrates in tumors will be of particular interest [141].

Similarly, most macromolecular drugs (including nucleic acids and some proteins) cannot readily permeate through the cell membrane and reach their pharmacological target. For these classes of API the modification of the NPs with targeting ligands may be more appropriate. Active targeting increases the affinity of the NPs for tumor cells, increasing its tumor residence times and allowing the drug-loaded NPs to efficiently enter the cells through receptor mediated endocytosis. The principles and future development of active targeting using affinity ligands on the surface of NPs will be presented in the next section.

3. Active targeting: toward a magic bullet?

Active targeting, also called ligand-mediated targeting, involves utilizing affinity ligands on the surface of NPs for specific retention and uptake by the targeted disease cells. To that end, ligands are selected to bind surface molecules or receptors overexpressed in diseased organs, tissues, cells or subcellular domains [6,33,142–144]. Actively-targeted material needs to be in the proximity of their target to benefit from this increased affinity. Therefore, the approach is aimed toward increasing interactions between NPs and cells and enhancing internalization of drugs without altering the overall biodistribution [6,145,146].

The design of actively-targeted NP drug carriers is complex because the NP architecture, the ligand conjugation chemistry and the types of ligand available all contribute to the efficacy of the system. Other factors like the administration route or the non-specific binding of proteins during the NP's journey through the bloodstream have been shown to affect the targeting ability of NPs. [12,147]. Physicochemical properties like the ligand density [148], the size of the NPs [149] or the choice of the targeting ligand [150] might also possibly affect the efficacy of the active targeting strategy *in vitro* and, most importantly *in vivo*. The following section will highlight the strategies, benefits and drawbacks of combining targeting ligands with NP drug delivery systems in the targeting of solid tumors.

3.1. The fundamentals of active targeting

The main mechanism behind active targeting is the recognition of the ligand by its target substrate. Representative ligands include antibodies, proteins, peptides, nucleic acids, sugars, and small molecules such as vitamins [151]. Target molecules can be proteins, sugars or lipids present in diseased organs or on the surface of cells [152,153]. The interactions of ligand-functionalized NP systems with their target antigen are enhanced by the multivalent nature of the NP architecture: multiple copies of the ligand increase the avidity of the NP for its target [154]. The targeting specificity and the delivering capacity are two important aspects to evaluate the efficiency of an active targeting system. The specificity is determined by the biodistribution of the ligand-functionalized NP and by how the conjugated ligand and the NP system interact with off-target molecules and cells; it is defined by the ligand and NP properties. The delivering capacity is directly related to the NP material and structure [33,155]. Currently, actively targeted NPs are envisioned as a promising complementary strategy to EPR to further augment the efficiency of cancer nanomedicines.

Actively-targeted NPs require being in the vicinity of their target antigen to recognize and interact with it. That intrinsic characteristic is considered a major challenge to the development of actively-targeted

NPs [156]. Here again, the systemic clearance of the NPs affects the amounts available in the bloodstream supplying the tumor. Because tumor blood flow is low compared to those observed in the organs of the MPS [12], the increase in the NPs' affinity for the targeted tumor-antigens cannot always compensate for the natural clearance processes. Therefore, Actively-targeted NPs need to be designed to have extended blood circulation times. Similarly, because molecular targets are usually situated in the extravascular space of the tumor, NPs rely on the EPR effect to reach their targets [113,114]. Together, these factors explain why active targeting strategies cannot radically change the biodistribution profiles of nanomaterials [157–159] and why the blood circulation times of ligand-decorated NPs need to be optimized to achieve optimal targeting [4,33,160].

Active targeting has been efficiently exploited to increase the NP internalization by the target cells and improve the efficacy of their payloads [4,158,159,161]. Anti-HER2 targeting moieties on the surface of liposomes strongly increase the uptake of the NPs in HER2-expressing cancer cells [158]. In opposition, non-targeted liposomes or targeted liposomes injected to mice bearing non-HER2-expressing tumors were shown to accumulate in the perivascular and stromal space in higher proportions [158]. In those cases, the liposomes were highly captured by the macrophages and showed reduced interactions with the cancer cells. Similarly, the intracellular delivery of nucleic acids can also take advantage of active targeting. For example, Bartlett *et al.* showed that the targeting of the transferrin-receptor is essential to the silencing of a luciferase beacon in a neuroblastoma xenograft [159]. Evidently, NP intracellular trafficking is complex [162] and receptor-mediated internalization can qualitatively affect the vesicular transport [163,164]. Actively targeted NPs that increase therapeutic efficacy will circumvent these issues by being capable of sufficient endosomal escape or by encapsulating drugs that are impervious to the endosomal/lysosomal environments.

Recently *in vitro* works on targeting concomitantly multiple surface receptors with a single NP were conducted [165,166]. The presence of both ligands (ligands targeted to folic acid combined to either EGFR antibodies [165] or glucose [166]) seemed to quantitatively improve cellular internalization. *In vivo* evaluation of non-sterically stabilized polystyrene nanoparticles functionalized with antibodies (against transferrin and intercellular adhesion molecule-1) seemed to show that multiplexed ligands could affect biodistribution [167]. However, because the nanoparticles in this study are so remote from potential therapeutic NPs (in terms of size and hydrophobicity, for example), it remains difficult to assess if such approach will really prove beneficial to target cancer cells.

3.2. Ligand conjugation/attachment strategies

Since NP avidity is directly related to the ligand density, the approaches used to introduce ligands on the surface of the NP are key aspects of actively-targeted systems. In most cases, covalent attachment of the ligand is the preferred strategy, but physical adsorption using affinity complexes can also be used effectively [168].

Because Organic and inorganic materials have different physicochemical properties, the type of particle will determine the difficulty of the ligand conjugation step [169]. For example, the surface functionalization of gold surfaces can be carried by directly reacting thiols with the surface [170], but other inorganic materials need the introduction of functional groups (e.g. NH₂ and OH) to increase their reactivity [171–173]. Organic polymers require different strategies where side chains or terminal reactive functions are reacted before or after NP synthesis.

In all cases, the stability of the ligand–NP bond will dictate how the particle retains the targeting moiety on its surface. In that regard, other design considerations must also be taken into account. For example, polymer matrices that erode homogeneously in bulk [174,175] might be preferable to surface-eroding polymers [176] for which the tethered

ligand can shed as the polymer degrades. Similarly, ligands that are non-covalently inserted in a lipid bilayer might require bulkier hydrophobic anchors to remain stable *in vivo* [138,177]. The following section will discuss the ligand conjugation/adsorption strategies in detail, with a specific emphasis on commonly utilized biodegradable polymers [178].

3.2.1. Pre-conjugation vs. post-formulation strategies

The conjugation of the ligand to the NP material is relatively straightforward when it is done before the assembly of the NP formulation. Pre-conjugation can be achieved with small molecules [150], peptides [179] and aptamers [142]. It is less adapted to native proteins with complex secondary structures as the conjugation step usually involves exposure to organic solvents. The pre-conjugation enables a subsequent one-step formulation procedure that reduces the risks of side reactions, and allows greater control over NP properties. Furthermore, this strategy allows the introduction of multiple types of ligands to NPs and facilitates the purification steps after NP formulation.

The alternative to this pre-formulation strategy is post-formulation conjugation where the ligands are reacted with formulated NPs. This strategy works for all types of ligands: antibody, protein, peptide, aptamer and small molecules, and might be preferable when the stability of the ligand in organic solvents is an issue, when the size of the ligand is too large to promote self-assembly or when the presence of the ligand changes the physicochemical properties of the NP components. For example, some proteins lose their functions in organic solvents and they are too large to participate in the NP self-assembly process [180]. In other instances, hydrophobic molecules tethered at the end of flexible copolymer chains modify their solubility and the ligands end up embedded in the hydrophobic core during the NP formation [150]. In those cases, it can usually be beneficial to conjugate these ligands after NP formation, directly onto the NP surface.

3.2.2. Synthetic strategies for conjugation

Bifunctional linkers are used to conjugate ligands to polymers or NPs via a series of chemical coupling reactions. The formation of a peptidic bond between the ligand and the NP surface is usually done by activating carboxylic groups (using, for example, N-hydroxysuccinimide (NHS) and 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC)) and reacting it with nucleophilic groups (e.g., amine) on the ligand. Although this approach is straightforward and can be carried in both aqueous and organic environments, the selectivity of the conjugation and the final orientation of the ligand both depend on the number of nucleophilic functions on the ligand.

Similarly, another coupling strategy couples a maleimide group on NP surfaces or polymers with a thiol group present on the ligand (i.e., cysteines in proteins or peptides) to form a stable thioether bond. In proteins or peptides, disulfide bonds can be reduced to free thiols to allow the reaction, provided that the reduction does not alter the three-dimensional structure and affinity of the ligand for its substrate. Similarly, free thiols can be introduced by first reacting 2-iminothiolane to primary amines, but this approach introduces a positive charge on the NP–ligand construct [181]. Other chemical efforts have also been reported for introducing thiols to molecules [182].

Recently, a bioconjugation method called “click chemistry” was developed which is a single step reaction that involves heteroatom bonds with or without catalysts [183]. For example, alkyne groups on peptides or small molecules will readily form a bond with the azide group on the NP surface or polymer backbone without side reactions. This method is very selective and provides very good yields. Copper-catalyzed click-chemistry is limited by the strenuous purification required to remove the toxic catalyst and by the fact that copper can have deleterious effects on the ligands themselves. For these reasons, non copper-catalyzed click-chemistry has been developed using cycloadditions such as cyclooctyne, tri(benzyltriazolylmethyl)amine, and sulfonated bathophenanthroline [183]. Finally, another limitation is that

alkyne and cyclooctyne groups are artificial chemical moieties and must be introduced synthetically to the ligand structure. When the ligand is a peptide, this can be easily done by introducing a reactive amino-acid at the C- or N-terminus. This strategy is less viable for small molecules where the new, relatively bulky functional groups can compromise affinity or for larger, more complex proteins produced through recombinant or bioengineering strategies.

3.2.3. Non-covalent approaches

Streptavidin–biotin interaction may be helpful to overcome limitations of synthetic bioconjugation [168]. Nanoparticles coated with avidins have complementary pairings with biotinylated ligands. Biotin is commonly used due to its small molecule size which does not alter the functions of the ligands [184]. This strategy is more versatile in comparison to bioconjugation method and has been applied to antibodies, peptides and aptamers [185]. It is particularly useful to establish proof of concepts or to screen different ligands without interference from the conjugation chemistry or to study fundamental ligand decoration parameters [185,186]. The weakness of this method is that it may cause immunogenicity due to the presence of the exogenous protein on the surface and it is not usually suitable for human use [187]. Also, because avidin can bind biotin in a multivalent fashion, cross-linking between the NP constituents is possible.

Alternatives to the avidin–biotin strategy use other high affinity complexes like Vancomycin/D-Ala–D-Ala [188] or adamantane–cyclodextrin coupling [189]. For example, the latter strategy allows the conjugation of ligand and adamantane to hydrophilic polymers (like PEG) using very mild conditions, and the subsequent tethering of the PEGylated-ligand on the surface of cyclodextrin-bearing nanoparticle via non-covalent interactions [190].

3.3. Influence of the architecture of actively-targeted NPs

The conjugation of ligands on the surface of NPs changes the properties of both the targeting molecules and the nanomaterial [33,142]. While ligands lose the rotational and translational freedom bestowed to free molecules, the new targeted entity achieves improved avidity because of the increased valency [149,154,191]. Similarly, the size, geometry, surface properties (charge and hydrophobicity), and composition of NPs can also be altered (Fig. 5). In some cases, NPs have shown benefits that go beyond the simple delivery of drug. For example, strands of nucleic acids immobilized on the surface of nanomaterials are more resistant to nuclease degradation [192,193]. To fully understand the properties of actively-targeted NPs, it is critical to determine how the physicochemical properties of the NPs affect the interactions with their targets.

3.3.1. The ligand density

Because increased valency allows cooperative effects, the density of the targeting molecules on the surface of NPs impacts their affinity for the substrate. Thermodynamically, the binding of a ligand to its substrate facilitates the subsequent binding of its neighbors [191,194]. Biologically, the multiple interactions of the NP with the cell membrane force the clustering and local concentration of receptors. This triggers the wrapping of the membrane and leads to internalization [195]. Together, these incidences impede the detachment of the NP from the cell surface and result in increased avidity.

This allows the use of multiple relatively low affinity ligands to efficiently bind targets with high avidity [196]. *In vitro*, this increasing ligand density usually results in improved cellular uptake [148,197]. However, this increase in affinity is not always linear. In some cases,




Fig. 5. The physicochemical properties of the ligand and the NP affect their blood circulation profiles, their biodistribution and their ability to be internalized by cancer cells.

the cooperative effect of the ligand can saturate and further increases in ligand density can have deleterious effects on cell binding [197,198]. This effect can be explained by improper orientation of the ligand, steric hindrance of neighboring molecules or competitive behaviors for the binding of the receptor. Similar negatively cooperative systems have been observed with folic acid-targeted micelles where the ligands are arranged in patchy clusters [199]. In these NPs, the display architecture of the targeting ligands influenced the extent of receptor-mediated tumor uptake in cancer cells both *in vitro* and *in vivo*. In other cases, the use of high densities of hydrophobic ligands can increase the macrophage uptake of the NPs without providing significant advantages in terms of receptor-mediated internalization [150].

Similar effects are seen *in vivo* where higher densities do not however always result in improved efficacy. The alteration of the NP surface to incorporate the ligands can modify the blood circulation and biodistribution profiles of the material. Using aptamers to target polymeric NPs to prostate cancer cells, Gu *et al.* showed that increasing the ligand density above 5% caused increased clearance of the NPs by the liver and the spleen and impeded the distribution to the tumor [148]. In fact, the increased clearance of targeted NPs by the MPS is critical and has been responsible for the demise of most actively targeted systems [200].

Finally, it has been observed that tumor selective antibodies could suffer from a binding-site barrier preventing their in-depth diffusion in the tumor [201–203]. This phenomenon is observed when high affinity molecules rapidly bind perivascular cells upon their extravasation to the tumor ($\uparrow R_i$ in Eq. (2)) [204–207]. The binding-site barrier also limits the in-depth diffusion of small molecular weight drugs with high affinity for tumor cells [208]. This effect has been recently detected with 25-nm NPs targeted to the epidermal growth factor which showed limited tumor penetration compared to their non-targeted counterpart [209]. Interestingly in that case, the binding-site barrier effect was not observed with larger NPs (*i.e.*, 60-nm particles), presumably because of differences in the rates of tumor penetration and/or cell internalization. Therefore, it seems probable that adequately tuning the ligand density on the NP surface could mitigate the binding-site barrier effect and translate into adequate retention time and maximal cellular uptake throughout the tumor.

3.3.2. The NP size and shape

The size and shape of the nanomaterial must be taken into consideration early in the design of targeted NPs. For spherical particles, smaller sizes represent higher curvatures which can be problematic for post-synthesis ligand functionalization. For example, when focusing on the adsorption of relatively rigid protein A on the surface of gold NPs, very small NPs (5 nm) can result in poor or uneven ligand surface densities [210]. In this specific study, this effect was not observed with larger NPs (*i.e.*, 15 nm).

In addition, the tethering of high molecular weight ligands (*e.g.*, antibodies, proteins, aptamers) on the surface of NPs increase their hydrodynamic radius beyond that of the unfunctionalized material [149]. This increase in size must always be considered in light of the possible size restrictions involved in tumor accumulation (see Section 2.4.1).

Besides the abovementioned effects, size can also affect cellular uptake. Using gold and silver NPs targeted with anti-HER2 antibodies, Jiang and colleagues showed *in vitro* that optimum cellular uptake in breast adenocarcinoma cells was obtained with a very narrow size range, *i.e.*, 25–50 nm (R_H with antibodies: 45 to 80 nm) [149]. Although, the avidity of the particle increased with size between 2 to 70 nm (R_H with antibodies: 13 to 100 nm), the authors explain that maximum uptake is a compromise between high avidity and optimal cell membrane wrapping around the NP. As these experiments were conducted in culture, these size considerations only took into account interactions with the surface of the cells and not convective or diffusive aspects of intratumoral transport. Additionally, it is not clear how these conclusions can be expanded to other systems, as NP-cell interactions

highly-depend on the physicochemical properties of each material [211,212].

In vivo, Lee and colleagues reported that the size of actively targeted NPs (25 vs. 60 nm) could affect the intracellular deposition [209]. In this case, although the intratumoral accumulation of smaller NPs was decreased due to shorter blood circulation times, the cytoplasmic and nuclear distribution of the 25-nm NPs was superior to that of the 60-nm colloids.

Besides the effect on circulation properties and tumor accumulation aforementioned [82,109], the shape of NP seems to influence the cell uptake kinetics and internalization pathways by modulating the interactions between the nanomaterial and the cell surface [211,213]. The internalization of non-spherical targeted NPs has recently been studied [214,215]. Barua *et al.* showed that, when compared to 200-nm diameter spheres, HER2-targeted 370×125 nm nanorods improved specific uptake by 1.6 fold while reducing non-specific internalization [215].

3.3.3. Surface and ligand charge

From a synthetic perspective, the charge of the unfunctionalized NP and that of the ligand can affect the conjugation yield and the spatial display of the ligand on the surface [216]. Repulsive or attractive forces between the surface of the NPs and the ligand can interfere with the conjugation [217,218] or affect the final ligand structure and conformation. A chemical spacer with reasonable length, such as PEG, can be helpful to reduce the effect, but might simultaneously complicate synthesis and increase the final particle size [197].

As discussed in Section 2.4.2., the final surface charge will affect the efficacy of the targeted NPs. Due to the interaction between cationic NPs and negatively charged cell membranes, positively-charged NPs show increased cellular binding and uptake, in a non-specific manner [219]. As most ligands are charged molecules, the NP surface charge is determined by the combinations of ligand densities, materials, and NP formulation strategies. Although recent work was carried out to address how charge density affects interactions of actively targeted NPs with cells [220] and how optimization of the ligand densities and NP charge can affect cellular uptake [221], it remains unclear what parameters offer the best tumor targeting *in vivo*.

3.3.4. Surface hydrophobicity

Besides surface charge, hydrophobicity can also affect the architecture of the ligand display [148]. This can have serious effects since most polymeric NPs have hydrophobic cores (*e.g.*, polyesters, polyamides) [222]. Valencia *et al.* showed that during the self-assembly of polymeric hydrophobic particles, folic acid, a model hydrophobic ligand, could remain trapped in the particle core without being properly displayed on the surface [150]. This resulted in a non-linear rise of NP affinity with increased feed-ratios of ligands in the formulation. Since this effect was not observed for a more hydrophilic ligand (the RGD peptide), this study highlighted the necessity of thorough physicochemical characterization of NPs after synthesis.

The final surface hydrophobicity of the NPs can also affect non-specific interactions with cells. On the one hand, actively targeted NPs without steric stabilization seem to lose their substrate-binding capacity when proteins adsorb on their surface [223]. Using silica NPs functionalized with transferrin, Salvati and colleagues showed that NPs lost their selectivity for cells expressing the transferrin-receptor upon plasma incubation. Although raising an important issue, these *in vitro* works are preliminary: these particles were not optimized for prolonged blood circulation times, and probably would have faced many other issues *in vivo*.

On the other hand, while PEG surface-functionalization can delay adsorption of opsonins and plasma proteins, the use of long or dense PEG chains can also prevent ligands from reaching their targets. This phenomenon has been demonstrated *in vitro* [197,224] and *in vivo* where the efficacy is dependent on both circulation times

and ligand–substrate interactions [225]. Since minimal PEG coverage seems to be required to maximize circulation times and tumor distribution, NPs that lose their PEG protection in the vicinity of cancer cells might provide an opportunity worth investigating [139].

3.4. Targeting ligands

Choosing the right type of ligand is critical to the efficiency of actively-targeted NPs. The first targeted systems centered mainly on the use of antibodies as targeting moieties because of their high specificity and wide availability [200]. Since then, other proteins, peptides, nucleic acid-based ligands and small molecules have all been described (Table 2). In the following section, the current understanding of the benefits and limitations of each system will be highlighted. Various issues of ligands that may affect ligand-mediated targeting will also be discussed, including ligand MW/size, ligand surface property, and ligand density on NP surfaces.

3.4.1. Antibodies and their fragments

An antibody (Ab, also called immunoglobulin (Ig)) is a large Y-shaped glycoprotein that can recognize the specific parts of a foreign target (an antigen). The dimeric functional region (hypervariable region (HVR), also called F(ab')₂ fragment) at the tip of the antibody can have very large numbers of slightly different structures (binding sites), offering the possibility to recognize a variety of antigens. The Fc fragment, at the base of the Y-shape, is much less variable and is responsible for the recognition of the protein by the MPS and the immune system.

The interest in using Ab as targeting moieties stems from the important role they play in modern therapeutics. Since the 1980s, antibodies have been the most widely investigated targeting ligands in the clinic and more than 30 types of monoclonal antibodies (mAbs) have been approved for clinical use, including rituximab, trastuzumab, cetuximab and bevacizumab [30–33]. The conjugation of Abs on the NP surfaces

aspires to combine their specificity and affinity with the unique properties of the NPs themselves. The first reported examples of targeted NPs are liposomes decorated with mAbs [30–32]. Although these systems showed improved cellular uptake *in vitro*, antibody-targeted NP delivery systems still face many limitations and challenges, which greatly limit their pertinence *in vivo*.

First, antibodies are large proteins with a molecular weight of 150 kDa and a hydrodynamic radius of 15–20 nm. This large size impedes the effective surface conjugation and causes notable increases in the diameter of the NPs [159,244]. To circumvent this problem, smaller fragments of Abs were proposed as targeting moieties, but the monomers and dimers of the Fab recognition patterns still represent bulky molecules with sizes around 50 and 100 kDa, respectively [6,200].

Second, the physiological role of antibodies is to clear antigens from the circulation by facilitating their recognition by immune cells and the MPS. The conjugation of Abs on a nanosized carrier therefore results in very effective clearance from the blood [245,246]. Removing the Fc fragment (*i.e.*, using only Fab fragments) resulted in slight improvements of blood exposure, but still compromised circulation times compared to undecorated NPs [247]. Others have proposed different conjugation strategies using the Abs carbohydrates to proper position them on the NPs and minimize the display of the Fc segment on the surface of the particle [248], but the results of this novel preparation procedure had mitigated effects on the circulation times [249].

A third limitation of using Ab-decorated NPs is the relative sensitivity of these proteins to environmental challenges (temperature, salt concentration and enzyme) and their low resistance to organic solvents. This creates technical challenges for the reproducible scale-up of NP preparation, affects the cost/efficiency ratio of the preparation, and restricts their stability and shelf-lives.

Despite these challenges, a few Ab-targeted NPs have made it to the clinical stages; MCC-465 and SGT-53 are two examples of antibody fragment-targeted NPs (Table 3). MCC-465 showed positive results in preclinical studies with adequate biodistribution and highly efficient delivery of doxorubicin to stomach cancer cells [250]. Another example of an NP with a single chain antibody fragment is SGT-53, which targets transferrin-receptors (Tf-R) on tumor cell surfaces by targeted delivery of the p53 suppressor protein [251]. With evident tumor growth inhibition in multiple cancers (including head and neck, prostate, and breast), this platform has great potential for future clinical trials [252]. Finally, a group of scientists at the University Hospital of Basel recently published their results of the clinical investigation of PEGylated DOX liposomes (Doxil®/Caelyx®) incorporating a F(ab') fragment of cetuximab (Erbix®/anti-EGFR mAb) in 29 patients with solid tumors [253]. Their liposomes, manufactured according to good manufacturing practices at the hospital's pharmacy, showed promising activity including 1 complete response in a patient with head and neck cancer.

3.4.2. Other proteins

The three-dimensional shape of proteins provides affinity for specific substrates, and therefore non-antibody proteins can be used as targeting moieties. Numerous naturally-occurring proteins have endogenous targets that can be exploited for therapeutic applications. For example, transferrin (Tf) is a 80-kDa glycoprotein which is one of the most abundantly studied targeting ligands [189,254,255]. Physiologically, Tf is responsible for the transport and regulation of iron concentration in biological environments. On the surface of cells, it binds the internalizing transferrin-receptor (Tf-R) with high affinity. Because Tf-R is up-regulated on the surface of cancer cells, NPs decorated with Tf have attracted much attention for the delivery of anticancer therapeutics [256]. The presence of Tf-ligands was shown to be essential for the intracellular delivery and gene silencing efficacy of siRNA nanocomplexes [159,189]. The strategy of using Tf to target Tf-R is currently under clinical investigation for various NPs [143].

Synthetic proteins can also be exploited as targeting ligands. For example, affibodies [230] or ankyrin repeat proteins [229] were developed

Table 2

Various examples of targeting ligands used in preclinical studies. EpCAM: epithelial cell adhesion molecule; EDB: extra-domain B.

Ligand type	System	Target	Indication	Ref
<i>Antibodies and fragments</i>				
Full antibody	Quantum dots	PSMA	Cancer (imaging)	[226]
F(ab') ₂	Liposome	GAH	Gastric cancer	[227]
F(ab')	Liposome	HER2	Breast cancer	[228]
scFv	Liposome	HER2	Breast cancer	[158]
<i>Proteins</i>				
Transferrin	Polymeric NPs	Tf receptor	Cancer	[159]
Ankyrin repeat protein	siRNA complexes	EpCAM	Cancer	[229]
Affibodies	Polymeric NPs	HER2	Breast cancer	[230]
<i>Peptides</i>				
CGNKRTRGC (LyP-1)	Protein NPs	gC1qR (p32)	Cancer	[231]
F3 peptide	Iron oxide NPs	Nucleolin	Cancer (imaging)	[232]
iRGD	Iron oxide NPs	αvβ3/5	Cancer (imaging)	[233]
iRGD	Polymeric NPs	αvβ3/5	Cancer	[234]
KLWVLPKGGGC	Polymeric NPs	Collagen IV	Inflammation	[235]
KLWVLPK	Polymeric NPs	Collagen IV	Vascular wall	[236,237]
Aptides	Liposomes	Fibronectin	Cancer	[238]
<i>Nucleic acid-based ligands</i>				
A10 aptamer	Polymeric NPs	PSMA	Prostate cancer	[161,239]
A9 CGA aptamer	Gold NPs	PSMA	Prostate cancer	[240]
<i>Small molecules</i>				
Folic acid	Gold nanorods	FA receptor	Cancer	[241]
Folic acid	Polymeric NPs	FA receptor	Cancer	[242]
TPP	Polymeric NPs	Mitochondria	Various	[243]
ACUPA	Polymeric NPs	PSMA	Cancer	[4]

Table 3

Actively-targeted NPs currently in clinical trials. When possible, usual doses for the non-targeted drug payloads are given for comparison. CR: complete response; PR: partial response; SD: stable disease; PD: progressive disease.

Name	System	Target (ligand)	Active (usual dose)	Doses (recommended)	Available phase I/II results
BIND-014	Polymeric NPs	PSMA (small molecule)	Docetaxel (75 mg/m ²)	15–75 mg/m ² (60 mg/m ²)	28 patients [4]: 1 CR (cervical cancer); 3 PR (NSLC, prostate, ampullary); 5 SD for >12 weeks (pancreatic, colorectal, gall bladder, tonsillar, anal)
anti-EGFR-ILs-dox	Liposomes (Doxil®/ Caelyx®)	EGFR (Ab, cetuximab)	Doxorubicin (50–60 mg/m ²)	5–60 mg/m ² (50 mg/m ²)	29 patients (26 evaluable) [253]: 1 CR (larynx); 1 PR (mesothelioma); 10 SD for 8–48 weeks
MM-302	Liposomes	Her2 (Ab scFv)	Doxorubicin (50–60 mg/m ²)	8–50 mg/m ²	34 breast cancer patients (22 evaluable) [312]: 2 PR (1 with 100% regression of lesion); 12 SD
MCC-465	Liposomes	Gastric cancer (Ab fragment)	Doxorubicin (50–60 mg/m ²)	6.5–45.5 mg/m ² (32.5 mg/m ²)	23 recurrent or metastatic stomach cancer patients (18 evaluable) [309]: 10 SD for 6–20 weeks
MBP-426	Liposomes	Tf-Receptor (Tf)	Oxaliplatin (75–85 mg/m ²)	6–400 mg/m ² (226 mg/m ²)	39 patients: 15 SD for 6 weeks; 3 SD for 12–24 weeks (colorectal)
SGT53	Liposomes	Tf-Receptor (Ab scFv)	p53 plasmid DNA (N/A)	0.6–3.6 mg (2.4 mg)	11 patients (10 evaluable)[310]: 7 SD at 6 weeks 3/3 patients with transgene plasmid detectable in tumor
CALAA-01	Polymeric NPs	Tf-Receptor (Tf)	siRNA M2 subunit of ribonucleotide reductase (N/A)	18–30 mg/m ²	3 patients [2]: 3/3 patients with decreased mRNA expression in tumor
Rexin-G	Retroviral vector	collagen (viral envelope peptide)	Human cyclin G1 gene (N/A)	1–6 × 10 ¹¹ CFU per 1-week cycle [315] 8–24 × 10 ¹¹ CFU per 4-week cycle [313,314]	12 metastatic pancreatic cancer patients [315]: 1 SD for 4 weeks (symptomatic deterioration); 11 PD for 4–7 weeks 13 chemotherapy-resistant pancreatic cancer patients (9 evaluable) [313]: 8 SD for 12–30 weeks; 1 PR for 52 weeks 42 metastatic sarcoma and osteosarcoma patients (37 evaluable) [314]: 23 SD for 5–15 weeks

to decorate NPs. This approach possesses the advantage of using high affinity, artificial ligands which do not have to compete against highly abundant, naturally-occurring proteins. Peptide aptamers are also synthetic fusion proteins where short variable peptide domains are confined by a constant protein scaffold [257]. This unique double-constrained structure over a loop of 10–12 amino acids offers binding affinities similar to those of antibodies. The protein scaffold may be any soluble compact protein, such as bacterial Thioredoxin-A [258]. Peptide aptamer selection can be made from combinatorial peptide libraries constructed by phage display and other surface display technologies such as mRNA, ribosome, bacterial and yeast displays [259].

Protein ligands share some limitations with antibodies: their bulky nature results in significant increases in size and their patterning on the surface of NPs can trigger immune responses. In fact, the conjugation of proteins on polymeric substrates can affect their metabolism and elimination pathways [260,261]. Also, because the amino acid sequences are generally more complex, the conjugation procedures are less straightforward. The many reactive functional groups in their structure (*i.e.*, amine groups or cysteines) complicate their optimal arrangement on the NP surface and can lead to cross-linking between NPs. Also, because their tertiary structure is usually essential to their affinity for their antigen, proper precautions must be used to ascertain that the conjugation processes do not abolish their ligand properties.

3.4.3. Peptides

Peptides are linear or cyclic sequences of amino acids. They are typically differentiated from their larger counterparts (*i.e.*, proteins) by having sequences limited to less than about 50 residues. These shorter chains lead to smaller molecular sizes and simpler three-dimensional structures which result in improved stability and resistance to the

environment as well as easier synthesis and conjugation. Their smaller sizes also allow the use of pre-formulation conjugation techniques for the preparation of NP systems. These advantages, combined with improved screening techniques to isolate ligand–substrate combinations have contributed to the increased role of peptides as targeting moieties in the past decade.

The most widely investigated peptide ligand is probably the RGD (arginine–glycine–aspartic acid) peptide family, which can strongly and specifically bind to $\alpha v \beta 3$ integrin receptors [33,142]. Many peptide containing RGD sequences have been developed to target both cancer and angiogenic endothelial cells expressing the $\alpha v \beta 3$ receptors. One example of RGD is the cyclic peptide cyclo(-RGDfV-) (Cilengitide) used as an anti-angiogenic agent [262,263]. Despite the widespread applications, the nonspecific targeting of RGD restricts its drug delivery applications because $\alpha v \beta 3$ integrin is also widely expressed on normal or inflamed tissues. Some of the intrinsic properties of RGD may also affect its targeting efficiency, such as molecular geometry. Colombo *et al.* investigated the structure–activity relationship of linear and cyclic RGD peptides. In their hands, the cyclic ligand showed more than 10-fold higher anti-tumor efficiency than its linear counterpart [264]. In parallel, other new generations of RGD analogs with improved targeting potential are underway [33].

For reasons explained earlier, vasculature targeting is an interesting complement to the EPR effect. Endothelial cell-penetrating peptides can facilitate the transport of NPs across the cell membrane and into specific cell organelles. Some of these penetrating peptides contain a specific C-terminal C-end Rule (CendR) sequence [233], (R/K)XX(R/K), which is responsible for the penetration activity. These penetrating peptides (*e.g.* LyP-1 [265], iRGD [233], and F3 [266] peptide sequences) can interact with neuropilin-1 to facilitate the cell and tissue penetration

of NPs [267]. Roth *et al.* showed that the conjugation of the cyclic peptide LyP-1 (sequence: CGNKRTRGC) on NPs can penetrate through blood vessels [267]. Sugahara *et al.* also reported that iRGD peptide (peptide sequence: CRGDGKPDG) could enhance the efficiency of drug delivery [233]. Luo *et al.* reported that the LyP-1-functionalized NPs target the lymphatic tumor cells with high binding affinity and increase cellular uptake of the NP by 8-folds compared to non-targeted NPs [268]. In addition, LyP-1 is found to be internalized by its targeted tumor cells indicating its cell-penetrating abilities [233,269]. Other cell penetrating peptides for targeted NPs include Cys-Arg-Glu-Lys-Ala (CREKA) [270], Asn-Gly-Arg (NGR) [271], and Ile-Thr-Asp-Gly-Glu-Ala-Thr-Asp-Ser-Gly (LABL) [272].

Finally, peptides can elicit agonist or antagonist pharmacologic activity on their substrate. This activity can alter the fate of NPs decorated with such peptides [273]. In this study, the targeting of quantum dots to G protein-coupled receptors using peptide agonists led to the internalization of the nanomaterial. On the other hand, the use of a small molecule antagonist as a ligand commanded the specific binding of the NPs without cellular uptake. Presumably, this concept could be further exploited to more closely control the fate of targeted NPs.

3.4.4. Nucleic acid based ligands

Nucleic acid-based aptamers are another class of ligands with completely different structures from peptide-based aptamers. Nucleic acid aptamers (Apts) are single-stranded (usually short) oligonucleotides, such as DNA, RNA or modified nucleic acids (xeno-nucleic acids or XNA) [259,274]. Thanks to their unique conformational structures that originate from intramolecular Watson–Crick interactions, Apts show high affinity and specificity. Candidates are screened from large oligonucleotide libraries with random sequences by taking advantage of the nucleic acid sequences. Binders are selected and specifically amplified at the detriment of non-binders using the polymerase chain reaction (PCR). Apts with strong binding characteristics for various targets, including small molecules [275] and proteins [276] have been isolated. Although the relative ease of isolating high affinity ligands against a variety of substrates remains the largest advantage of Apts, their reproducible synthesis and the simplicity of their chemical derivation are also among the other benefits [259].

Although Apts can exist naturally in riboswitches [277–279], most of the Apts used for the design and synthesis of targeted NPs are artificially-engineered. However, although the diversity of possible sequences offers virtually unlimited binding options, the key element remains in the isolation of successful binders from non-binding candidates. Since the 1990s, the isolation of ligands has gone from simple chromatographic separation [280–282], to cell-based screening methods [283,284] and *in vivo* selection [285]. These processes evolved with the objective of finding candidates for increasingly complex substrates and specific applications. When screening for ligands intended for the design of actively-targeted NPs, the internalization of the target remains an important benchmark. Recently, a method was developed to exclude non-internalizing ligands from the enrichment procedure [284].

In vitro feasibility of Apt-decorated NPs to target the prostate-specific membrane antigen (PSMA) cancer marker was reported in 2004 [286]. Since then, the proof of concept was achieved *in vivo* for NPs incorporating multiple types of drug [160,161,241,287,288]. Overall, the targeting of PSMA showed improved efficacy in decreasing tumor growth and increasing survival. For example, Bagalkot *et al.* reported that the physical conjugate of Apt-Dox could result in almost 2-folds increase in survival rates in comparison to non-targeted NPs [287]. Similarly, intratumor injections of PSMA-targeted NPs were shown to delay tumor growth for more than 3.5 months [161].

Despite their added promises at the preclinical stage, the clinical development of Apt-decorated NPs has been impeded by certain limitations. First, because nucleic acids in biological environments are easily degraded by nucleases, the stability of the ligand on the surface of the

NPs is a concern. Although many strategies have been employed to prevent the degradation and excretion of free therapeutic Apts in the bloodstream [289], not much work has been conducted to specifically address degradation of Apts on the surface of NPs. By analogy, the conjugation of the Apt on the surface of NPs could provide steric protection similar to that conferred by PEG, sugars and cholesterol modifications [289]. Similarly, the use of nucleic acids modified with fluoro, amino, or methoxy groups [259] or artificially-locked nucleic acids (LNA) [289] should also delay degradation by nucleases.

Another concern for the surface functionalization of NPs with Apts is the effect that these ligands have on the circulation kinetics of the nanomaterial. Given the phosphodiester backbone of the nucleic acid chain, Apts have negative charges which can affect their circulation times [12]. Gu *et al.* showed that Apt densities above 5% resulted in decreased distribution to the tumor and increased capture by the organs of the MPS 24 h after intravenous injection [148].

Besides these fundamental considerations, other technical aspects of using Apts to decorate NPs are also important. Like other macromolecular ligands, Apts can increase the hydrodynamic radius of NPs. In fact, despite the fact that their size is relatively small (around 10–20 kDa), the stiff chain structure of the nucleic acid chains makes their size (length) similar to that of proteins (*i.e.*, from 3 to 10 nm) [259,290]. Furthermore, although the production costs have fallen recently due to technological developments and the expiration of patents, the selection and large-scale production of Apts on solid phase supports remain expensive procedures [192]. The production costs might be partially mitigated by secondary selection techniques which delete or shorten the non-critical regions while keeping the functional binding domains [259].

3.4.5. Small molecules

Small molecular weight compounds have properties which strongly contrast from the targeting ligands presented above: small sizes, low production costs, and improved stability. These advantages translate into simple pre-formulation conjugation strategies and simple, tunable NP synthesis. Hrkach *et al.* recently reported the preclinical developments leading to the clinical evaluation of PSMA-targeted NPs in an article that highlights many of the benefits of using small molecules as targeting agents [4].

In fact, the main challenge with small molecules is the identification of new affinity ligands for the substrates of interest. In general, the screening of small molecules is difficult to multiplex because selection and signal intensifying mechanisms like bacteriophage-mediated replication or PCR amplification do not exist. Notwithstanding a few examples of DNA-encoded libraries that facilitate the screening of binders using nucleic acid footprints [291,292], the identification of small molecular weight ligands involves serendipity or tedious high throughput screening procedures. For that reason, clinically-relevant ligand/substrate pairs are scarce and most of the examples of actively-targeted NPs rely on using widely known ligands.

Among the synthetic and natural small molecule ligands reported, a common example is folic acid (*i.e.*, folates or vitamin B9). Folates have been widely utilized because of their very high affinity ($K_D = \sim 10^{-9}$ M) and specificity for folate receptors (FR) which are frequently over-expressed on the surface of a variety of human tumors, including ovarian, brain, breast, colon, renal and lung cancers [293,294]. For example, folate-targeted NPs co-encapsulating paclitaxel and yttrium-90 showed improved survival in a xenograft model of ovarian cancer [242] and folate-targeted fluorescent dyes were investigated in humans to improve surgical debulking of tumors using intraoperative tumor-specific visualization [295]. On the downside, FR expression seems to be patient-dependent and must be assessed individually for each type of cancer [33]. Recently, folate-targeted imaging modalities were proposed to identify FR-positive patients [296]. In a personalized medicine setting, such an agent would help screen *a priori* the patients that could benefit from folate-

targeted therapeutic NPs. More importantly, as opposed to methods based on the *ex vivo* quantification of FR in lysed patient biopsies [297], *in vivo* imaging modalities might enable the quantification of FR on the surface of cancer cells, actually available for interactions with folate-decorated NPs. This is particularly valuable as it has been observed that as much as 50–75% of the FR pool can be localized in the endosomal membranes [294,298]. Finally, another limitation of folate-mediated cancer nanomedicines is that FR is constitutively-expressed in healthy tissues and normal epithelia of many organs, potentially limiting the selectivity of the ligand for diseased cells.

Another example of small molecule targeting ligands is triphenyl-phosphonium (TPP) and its derivatives for mitochondria targeting [243,299]. TPP is a cationic, relatively large and hydrophobic molecule that can penetrate easily through the cell membrane. An investigation indicated that the positively charged TPP could accumulate several hundred folds within mitochondria [299]. An *in vitro* study recently established the mitochondria targeting potential of TPP-decorated NPs [243]. These studies showed that NPs could efficiently be internalized and escape the endosomal compartment when their size was around 100 nm and their zeta-potential was higher than +22 mV. Although this positive charge, conferred in part by TPP, seems important for *in vitro* performance, it might possibly undermine its blood exposure *in vivo*. This factor will have to be considered in the future developments of this targeting strategy.

Carbohydrate moieties [300], including mannose [301], glucose [166], galactose [302], and their derivatives, have also been widely utilized as targeting ligands. Carbohydrates are recognized by ubiquitous cellular membrane proteins called lectins [303,304]. In fact, one of the first actively-targeted anticancer nanomedicines evaluated in humans was a DOX-conjugate, which incorporated a galactosamine targeting moieties to target primary and metastatic liver cancers [305]. Although this product failed in phase II due to lack of efficacy [306], it was observed in humans that the ligand conferred some level of liver targeting to this polymer–drug conjugate [305].

To identify other ligands, Weissleder and colleagues screened 146 different small molecules to act as targeting ligands [196]. Using highly reactive small molecules and magnetic particles to facilitate large-scale synthesis and purification, they screened their multivalent particles for cellular uptake in 5 different cell lines. The multivalency of 2 low affinity ligands on the surface of the NPs resulted in specific and efficient tumor targeting *in vivo*, after intravenous administration.

Finally, one of the most successful stories of actively-targeted NPs to date is based on the use of S,S-2-[3-[5-amino-1-carboxypentyl]-ureido]-pentanedioic acid (ACUPA), a small molecule targeted to PSMA [4,142]. ACUPA was initially identified as part of a prostate cancer specific imaging agent [307]. The decoration of the surface of docetaxel-loaded NPs with ~200 molecules of ACUPA resulted in optimal targeting of PSMA-positive prostate cancers without interfering with blood circulation kinetics [4]. The PSMA-specific, actively-targeted NPs, BIND-014 has now promisingly completed its phase I evaluation in humans and is currently under further clinical investigation.

3.5. Active targeting in humans

Although no actively-targeted NPs are currently commercially-available, at least 5 targeted liposomes and 2 targeted polymeric nanoparticle therapeutics have made it to clinical development stages. Among the NPs studies MBP-426 [308], MCC-465 [309], SGT53 [310], MM-302 [311,312], BIND-014 [4], CALAA-01 [2], cetuximad-decorated Doxil®/Caelyx® liposomes [253] and a retroviral vector [313–315] have results of phase I/II clinical trials available (Table 3). More details about these NPs and others ligand-functionalized therapeutics are examined in detail in a very good review article published during the preparation of this manuscript [316]. Fig. 6 also provides some examples of promising technologies currently studied in patients. The main therapeutic targets of these NPs are the Tf-R [2,143], the epidermal growth factor receptor [253,317], PSMA [4], the surface of gastric cancer cells [309] and the human epidermal growth factor receptor 2 (HER-2)




Fig. 6. Actively-targeted NPs have shown promises in early clinical trials. A. BIND-014, a PSMA targeted, docetaxel-containing polymeric NP has shown impressive anticancer response in heavily-pretreated patients; the regression of lung metastases experienced by one patient suffering from cholangiocarcinoma after two cycles of BIND-014 is evidenced here by CT scans. B. SGT53, TfR-targeted liposomes containing plasmid DNA for the p53 gene were shown to allow expression of the exogenous p53 gene by DNA PCR; tumor biopsies were taken 100 (T1) and 26 h (T2) after administration of different doses in 2 different patients. C. CALAA-01, a TfR-targeted polymeric NPs encapsulating siRNA was shown to reduce mRNA (77%) and protein expression (32%) compared to baseline (C2_{pre}) in one patient receiving 30 mg/m² of siRNA (C2_{post}); mRNA expression was assessed by qRT-PCR while protein expression was evidenced by western blotting. Figures are used with permissions from [2,4,310], respectively.

[312]. In the next few years, as the evaluation of these NPs in patients continues, our understanding of the parameters governing active targeting will hopefully improve. Nevertheless, a few important considerations regarding the use of actively-targeted NPs in humans must be highlighted.

3.5.1. The choice of the target

The design of actively-targeted NPs for clinical use involves scientific and economic considerations alike. Evidently, all important tumor biology and NP design aspects discussed earlier must be accounted for in order to develop the most efficient product possible. However, because of the large amount of investment involved in the development of therapeutic products, the probability of success, the prospective therapeutic impact and many other economic factors must be evaluated before choosing the target of interest [143].

The development of targeted drug delivery platforms is often considered to be more costly than the development of non-targeted NPs [143]. This belief likely stems from the increased complexities associated with the preparation and manufacturing of industrial-scale, clinical-grade NPs which display targeting ligands. That being said, the financial burden of scaling up and manufacturing must be considered in the development of any type of therapeutic; while it represents a significant proportion of the expenses associated with early development, the costs associated with large-scale clinical trials are certainly much more prohibitive. Once the engineering obstacles are surpassed and NPs show clinical efficacy, targeted NP platforms present unique designs which possibly confer long-term commercial advantages [318]. All things considered, the major challenge in the development of targeted and untargeted NPs alike remains to show sufficient efficacy in phase III clinical trials to warrant commercialization. If the presence of the targeting ligand confers better clinical efficacy, the risk and expenses related to more complex chemistry, manufacturing and control (CMC) procedures might potentially be compensated when the NPs reach commercialization.

Additionally, the population that would eventually benefit from the treatment must also be considered carefully. Given the wide range of cancer types in which the Tf-R is upregulated [319], it is not fortuitous that many NPs currently under study target this specific surface protein [33]. The prospect of having one versatile targeted platform that could simultaneously treat multiple cancers is very appealing to clinicians and investors alike. The shortcoming of this approach however, might emanate from the use of proteins as the targeting ligands; large and complex molecules can cause CMC issues that might hinder or limit the large-scale production and commercialization of the NPs. Furthermore, the targeting of ubiquitously-expressed proteins increases the risks of non-specific distribution and off-target toxicities. Poor tolerability and dose-limiting side-effects restrain the therapeutic regimens that can be utilized in patients. Such limitations have been the demise of many drug delivery systems (both targeted and non-targeted), when doses could not be titrated high enough to achieve therapeutic efficacy [320–322]. Even for commercially-approved, non-targeted nanomedicine, patients experience unique toxicities associated with the encapsulated form of the drug (e.g., hand–foot syndrome or complement activation-related pseudoallergic reactions with PEGylated liposomal DOX [323,324]). Since the clinical experience with targeted NPs remains limited, the appearance in clinical trials of non-expected toxicities due to the targeting ligand can never be completely ruled-out. As such, improved tolerability of anticancer therapeutics should therefore remain one of the main objectives throughout development.

In contrast, using molecular targets which are specific to a small population raises the hallmarks of required performance. When the number of prospective patients is small, a drug needs to be paradigm-shifting and become an unavoidable part of this population's therapeutic regimen to achieve good return on investment. The relative commercial successes of some small molecules targeted to very specific cancer metabolic pathways exemplify this phenomenon (e.g., crizotinib and

the ALK mutation in 3–5% of NSCLC patients). On the other hand, a drug that provides only marginal benefits might be overlooked and diluted by other comparable treatments. Such problems were encountered by SMANCS, a treatment mainly intended for hepatocellular carcinoma [70], and by others [305] for which the interest dwindled as clinical hurdles were exposed.

A good compromise is using targets like PSMA which are not constitutively present in healthy tissues, but expressed in the neovasculature of multiple types of cancer. Despite its name which suggests specificity for prostate cancer, PSMA is found in the vasculature of almost all types of solid tumors [325]. From an economic perspective, the versatility of the target therefore opens the door to multiple therapeutic indications and potentially higher return on investment. This is particularly interesting in the context of developing a high risk drug delivery platform. It also explains why, despite the fact that almost all of the preclinical data leading to PSMA-targeted NPs was obtained in prostate cancer models, the initial clinical investigation enrolled patients suffering from all types of cancer [4]. The results of the Phase 1 clinical trial in 28 patients showed one complete response (cervical cancer), 3 partial responses (prostate, non-small cell lung carcinoma (NSCLC) and ampullary cancer) and 5 stable diseases (pancreatic, colorectal, gall bladder, tonsillar and anal cancers), demonstrating that the targeted therapy could be beneficial in a variety of different solid tumors. These promising results offer the opportunity to continue further clinical investigations by stratifying patients in multiple types of cancer. In the case of BIND-014, phase 2 clinical trials were recently initiated, simultaneously in 3 independent studies: for the 1st line treatment of chemotherapy-naive, castrate-resistant prostate cancer as well as for 2nd line treatment of docetaxel-naive NSCLC and bladder cancer.

3.5.2. Assessing the impact of active targeting

Measuring the impact of active targeting in humans is not simple. In preclinical models, each individual component can be evaluated distinctively and in an iterative manner by testing different NPs. In humans, assessing the efficacy of a treatment requires time and resources; the parallel evaluation of matching targeted and non-targeted NPs remains therefore unlikely.

One way of addressing the problem if accurate phenotyping techniques are available [326] is by comparing patient cohorts which differ in the expression of the target biomarker. While this certainly raises the complexity and the costs associated with a clinical study, it also does not provide any significant advantages in the early evaluation of the NPs. In fact, at the critical step of assessing the clinical efficacy of a NP, it may be more rational to evaluate efficacy in homogenous, target-positive populations if such patients can be identified. Especially since in essence, the main criteria leading to failure or success of a drug is not the mechanistic aspects behind its performance, but how well it compares to standard treatments.

It is likely that the impact of active targeting might only be assessed in post-approval, phase 4 studies. Products are much less vulnerable to the outcome of mechanistic studies when they are commercially-available. This has been observed for approved non-targeted nanomedicines: for example, in retrospective investigations establishing correlations between therapeutic response and biomarker expression [327] or in prospective pharmacokinetic studies trying to predict the performances of nanomedicine [64]. Since ligand-functionalized NPs rely on sufficient surface expression of specific markers to exert their effect, resistance could theoretically arise from the natural selection of target-negative cell populations less exposed to the drug. Nevertheless, there is currently no evidence suggesting that such resistance would appear at a faster rate for targeted NPs than for any other anticancer agent. In fact, in certain instances, actively-targeted NPs were shown to overcome multidrug resistance, at least *in vitro* [316]. Mechanistic studies conducted once actively-targeted NPs are commercialized and used to treat larger patient cohorts should provide some understanding regarding potential

resistance mechanisms and how to counter them. More importantly, a better understanding of how these nanomedicines work will also provide insights on how to maximize the benefits for patients.

4. The near future of cancer nanomedicines

Since the EPR effect was first observed more than 30 years ago, the prospects for therapeutic NPs have never been brighter: the number of NPs in clinical trial is on the rise (examples of which are presented in Table 3), partnerships between pharmaceutical and nanotechnology companies are blossoming and our understanding of the interactions between nanomaterials and biological systems is increasing at an exponential rate. An impressive body of scientific evidences suggest that therapeutic NPs could shift the paradigm of treatment for numerous human diseases, including cancer, and that some NPs currently in development might become tomorrow's blockbusters.

Although most of the past efforts were focused on the development of differentiated drug carriers loaded with already-approved APIs, the true potential of therapeutic NPs might really materialize by emphasizing on otherwise “undruggable” compounds. Some molecularly targeted molecules necessitate precise pharmacokinetics and on-target activity to exert efficacy; while other drugs might benefit from improved biodistribution to mitigate toxicities and/or maintain therapeutic levels at the site of interest. Such difficult APIs might require the use of the NPs to ever become clinically-relevant; in which case the therapeutic value of the NP-API entity would result from the combination of drug and carrier, in an indissociable manner.

In the field of cancer more specifically, the complexity of the disease might compel the joint efforts of multidisciplinary teams. Material scientists may devise increasingly more robust methods for the design, synthesis and characterization of NPs. These will be important to improve our understanding of the interactions of nanomaterials with biological environments and to mitigate CMC challenges during development toward commercialization. From the cancer biology perspective, it is now increasingly clear that tumors are genetically heterogeneous and complex, and show variations in the tumor microenvironment and in many of the parameters postulated to be important in EPR. Likewise, from the clinical standpoint, inter- and inpatient genetic and phenotypic heterogeneities inherently affect drug response rates and NP distribution because of mutations in the diseased cells or alterations in the tumor microenvironment and vasculature. Patient enrichment might therefore be at the forefront of future therapies involving nanomedicines; to warrant optimal therapeutic efficacy, it might be crucial to identify *a priori* which patients will offer the best response to anticancer NPs. The hallmarks that highlight such predisposition might possibly be complex and encompass one or more different biomarkers. Streamlined assays to measure biomarker activity in the tumor during therapy might therefore be required. In parallel, imaging modalities developed as “companion diagnostic” may prove useful tools during treatment with NPs. For patients that do not exhibit the optimal phenotype warranting treatment with NPs, the perspective of artificially fine-tuning the tumor characteristics has been proposed by other investigators. Some clinical evidences of combinations of already approved drugs with NPs to augment the EPR exist (e.g., using angiotensin II to increase vascular pressure) [27].

Although the aforementioned elements hold true for the majority of therapeutic NPs, the relatively recent clinical evaluation of actively targeted systems offers an added degree of excitement in the field of cancer nanotechnologies. The community is eager to find out how well the evidences of improved drug exposure and retention in the tumor using actively-targeted systems in animals translate to humans. Several targeted nanoparticles are in clinical trials today, including the first targeted and controlled release polymeric NP (BIND-014), in Phase II [4]. As more comprehensive data sets become available, the clinical advantages of using actively targeted NPs might be better understood. Besides the benefits on tumor drug exposition cited earlier,

active targeting strategies might possibly prove useful to deliver APIs across physiological barriers, whether they are as simple as cell membranes, or as complex as the gastrointestinal epithelia or the blood brain barrier.

All together these elements forecast a bright future for non-targeted and targeted therapeutic NPs alike; in the next decade or so they might increasingly become a mainstay of safer and more efficacious treatments by ensuring proper localization of drugs at their site of interest. Although the processes of drug development are long, it is probable that nanotechnology will be able to impact medicine the same way it has influenced other scientific fields, like electronics, energy and material sciences. Such breakthrough might shift the paradigm of treatment for many diseases, including cancer, and contribute to a growing arsenal of non-targeted and targeted nanomedicines to tackle a myriad of important human conditions.

Acknowledgments

This work was supported by the National Cancer Institute (NCI) (grant U54-CA151884), the National Heart, Lung, and Blood Institute (NHLBI) Program of Excellence in Nanotechnology (PEN) (contract #HHSN268201000045C), the National Institute of Biomedical Imaging and Bioengineering (NIBIB) R01 grant (EB015419-01) and the David Koch-Prostate Cancer Foundation Award in Nanotherapeutics. Dr Nicolas Bertrand acknowledges a postdoctoral fellowship from the Canadian Institutes of Health Research (CIHR). Dr. Xiaoyang Xu acknowledges postdoctoral support from an NIH National Research Service Award (NRSA) (1F32CA168163-01). Dr. Farokhzad declares financial interests in BIND Therapeutics, Selecta Biosciences and Blend Therapeutics; three biotechnology companies developing nanoparticle technologies for medical applications. The rest of the authors declare no conflict of interest.

References

- [1] K. Strebhardt, A. Ullrich, Paul Ehrlich's magical bullet concept: 100 years of progress, *Nat. Rev. Cancer* 8 (2008) 473–480.
- [2] M.E. Davis, J.E. Zuckerman, C.H.J. Choi, D. Seligson, A.W. Tolcher, C.A. Alabi, Y. Yen, J.D. Heidel, A. Ribas, Evidence of RNAi in humans from systemically administered siRNA via targeted nanoparticles, *Nature (London)* 464 (2010) 1067–1070.
- [3] J. Tabernero, G.I. Shapiro, P.M. LoRusso, A. Cervantes, G.K. Schwartz, G.J. Weiss, L. Paz-Ares, D.C. Cho, J.R. Infante, M. Alsina, M.M. Gounder, R. Falzone, J. Harrop, A.C.S. White, I. Toudjarska, D. Bumcrot, R.E. Meyers, G. Hinkle, N. Svrzikapa, R.M. Hutabarat, V.A. Clausen, J. Cehelsky, S.V. Nochur, C. Gamba-Vitalo, A.K. Vaishnav, D.W.Y. Sah, J.A. Gollob, H.A. Burris, First-in-humans trial of an RNA interference therapeutic targeting VEGF and KSP in cancer patients with liver involvement, *Cancer Discov.* 3 (2013) 406–417.
- [4] J. Hrkach, D. Von Hoff, M.M. Ali, E. Andrianova, J. Auer, T. Campbell, D. De Witt, M. Figa, M. Figueiredo, A. Horhota, S. Low, K. McDonnell, E. Peeke, B. Retnajaran, A. Sabnis, E. Schnipper, J.J. Song, Y.H. Song, J. Summa, D. Tompsett, G. Troiano, T. Van Geen Hoven, J. Wright, P. LoRusso, P.W. Kantoff, N.H. Bander, C. Sweeney, O.C. Farokhzad, R. Langer, S. Zale, Preclinical development and clinical translation of a PSMA-targeted docetaxel nanoparticle with a differentiated pharmacological profile, *Sci. Transl. Med.* 4 (2012) 128–139.
- [5] D.A. Scheinberg, C.H. Villa, F.E. Escorcia, M.R. McDevitt, *Concepts of the infinite armada: systemic cancer therapy using nanomaterials*, *Nat. Rev. Clin. Oncol.* 7 (2010) 266–276.
- [6] D. Peer, J.M. Karp, S. Hong, O.C. Farokhzad, R. Margalit, R. Langer, Nanocarriers as an emerging platform for cancer therapy, *Nat. Nanotechnol.* 2 (2007) 751–760.
- [7] B.Y.S. Kim, J.T. Rutka, W.C.W. Chan, Nanomedicine, *New. Engl. J. Med.* 363 (2010) 2434–2443.
- [8] J.E. Kipp, The role of solid nanoparticle technology in the parenteral delivery of poorly water-soluble drugs, *Int. J. Pharm.* 284 (2004) 109–122.
- [9] L. Zhang, J.M. Chan, F.X. Gu, J.-W. Rhee, A.Z. Wang, A.F. Radovic-Moreno, F. Alexis, R. Langer, O.C. Farokhzad, Self-assembled lipid-polymer hybrid nanoparticles: a robust drug delivery platform, *ACS Nano* 2 (2008) 1696–1702.
- [10] K.A. Whitehead, R.S. Langer, D.G. Anderson, Knocking down barriers: advances in siRNA delivery, *Nat. Rev. Drug Discov.* 8 (2009) 129–138.
- [11] F. Alexis, E. Prigden, L.K. Molnar, O.C. Farokhzad, Factors affecting the clearance and biodistribution of polymeric nanoparticles, *Mol. Pharm.* 5 (2008) 505–515.
- [12] N. Bertrand, J.C. Leroux, The journey of a drug carrier in the body: an anatomo-physiological perspective, *J. Control. Release* 161 (2012) 152–163.
- [13] M. O'Brien, N. Wigler, M. Inbar, R. Rosso, E. Grischke, A. Santoro, R. Catane, D. Kieback, P. Tomczak, S. Ackland, F. Orlandi, L. Mellars, L. Alland, C. Tendler, Reduced cardiotoxicity and comparable efficacy in a phase III trial of PEGylated liposomal doxorubicin HCl (CAELYX™/Doxil(R)) versus conventional doxorubicin for first-line treatment of metastatic breast cancer, *Ann. Oncol.* 15 (2004) 440–449.

- [14] C.A. Geisberg, D.B. Sawyer, Mechanisms of anthracycline cardiotoxicity and strategies to decrease cardiac damage, *Curr. Hypertens. Rep.* 12 (2010) 404–410.
- [15] J. Cortes, C. Saura, Nanoparticle albumin-bound (nabTM)-paclitaxel: improving efficacy and tolerability by targeted drug delivery in metastatic breast cancer, *Eur. J. Cancer* 8 (2010) 1–10.
- [16] L.J. van 't Veer, R. Bernards, Enabling personalized cancer medicine through analysis of gene-expression patterns, *Nature* 452 (2008) 564–570.
- [17] S. Basu, P. Chaudhuri, S. Sengupta, Targeting oncogenic signaling pathways by exploiting nanotechnology, *Cell Cycle* 8 (2009) 3480–3487.
- [18] J. Baselga, Targeting tyrosine kinases in cancer: the second wave, *Science* 312 (2006) 1175–1178.
- [19] M. Martini, L. Vecchione, S. Siena, S. Tejpar, A. Bardelli, Targeted therapies: how personal should we go? *Nat. Rev. Clin. Oncol.* 9 (2012) 87–97.
- [20] J. Zhang, P.L. Yang, N.S. Gray, Targeting cancer with small molecule kinase inhibitors, *Nat. Rev. Cancer* 9 (2009) 28–39.
- [21] P.M. Valencia, E.M. Pridgen, B. Perea, S. Gadde, C. Sweeney, P.W. Kantoff, N.H. Bander, S.J. Lippard, R. Langer, R. Karnik, O.C. Farokhzad, Synergistic cytotoxicity of irinotecan and cisplatin in dual-drug targeted polymeric nanoparticles, *Nanomedicine* 8 (2012) 687–698.
- [22] Xiaoyang Xu, Kun Xie, Xue-Qing Zhang, Eric M. Pridgen, Ga Young Park, Danica S. Cui, Jinjun Shi, Jun Wu, Philip W. Kantoff, Stephen J. Lippard, Robert Langer, Graham C. Walker, Omid C. Farokhzad, Enhancing tumor cell response to chemotherapy through nanoparticle-mediated co-delivery of siRNA and cisplatin prodrug, *Proc. Natl. Acad. Sci. U S A* 110 (46) (2013) 18638–18643.
- [23] J.L. Yu, J.W. Rak, B.L. Coomber, D.J. Hicklin, R.S. Kerbel, Effect of p53 status on tumor response to antiangiogenic therapy, *Science* 295 (2002) 1526–1528.
- [24] Y. Sun, J. Campisi, C. Higano, T.M. Beer, P. Porter, I. Coleman, L. True, P.S. Nelson, Treatment-induced damage to the tumor microenvironment promotes prostate cancer therapy resistance through WNT16B, *Nat. Med.* 18 (2012) 1359–1368.
- [25] S. Sengupta, D. Eavarone, I. Capila, G. Zhao, N. Watson, T. Kiziltepe, R. Sasisekharan, Temporal targeting of tumour cells and neovasculature with a nanoscale delivery system, *Nature* 436 (2005) 568–572.
- [26] R.S. Tuma, Pancreatic cancer: gemcitabine plus nab-paclitaxel prolongs survival in patients with metastatic disease, *Oncol. Times* 35 (2013) 6–8.
- [27] U. Prabhakar, D.C. Blakey, H. Maeda, R.K. Jain, E.M. Sevick-Muraca, W. Zamboni, O.C. Farokhzad, S.T. Barry, A. Gabizon, P. Grodzinski, Challenges and key considerations of the enhanced permeability and retention effect (EPR) for nanomedicine drug delivery in oncology, *Cancer Res.* 73 (2013) 2412–2417.
- [28] P. Carmeliet, R.K. Jain, Principles and mechanisms of vessel normalization for cancer and other angiogenic diseases, *Nat. Rev. Drug Discov.* 10 (2011) 417–427.
- [29] L.D. Leserman, J.N. Weinstein, R. Blumenthal, W.D. Terry, Receptor-mediated endocytosis of antibody-opsonized liposomes by tumor cells, *Proc. Natl. Acad. Sci.* 77 (1980) 4089–4093.
- [30] L.D. Leserman, J. Barbet, F. Kourilsky, J.N. Weinstein, Targeting to cells of fluorescent liposomes covalently coupled with monoclonal antibody or protein, *Nature* 288 (1980) 602–604.
- [31] T.D. Heath, R.T. Fraley, D. Papahadjopoulos, Antibody targeting of liposomes: cell specificity obtained by conjugation of F(ab')₂ to vesicle, surface, *Science* 210 (1980) 539–541.
- [32] H.M. Warenius, G. Galfre, N.M. Bleehen, C. Milstein, Attempted targeting of a monoclonal-antibody in a human-tumor xenograft system, *Eur. J. Cancer Clin. Oncol.* 17 (1981) 1009–1015.
- [33] N. Kamaly, Z. Xiao, P.M. Valencia, A.F. Radovic-Moreno, O.C. Farokhzad, Targeted polymeric therapeutic nanoparticles: design, development and clinical translation, *Chem. Soc. Rev.* 41 (2012) 2971–3010.
- [34] N.M. Idris, M.K. Gnanasamandhan, J. Zhang, P.C. Ho, R. Mahendran, Y. Zhang, In vivo photodynamic therapy using upconversion nanoparticles as remote-controlled nanotransducers, *Nat. Med.* 18 (2012) 1580–1585.
- [35] H. Huang, S. Delikanli, H. Zeng, D.M. Ferkey, A. Pralle, Remote control of ion channels and neurons through magnetic-field heating of nanoparticles, *Nat. Nanotechnol.* 5 (2010) 602–606.
- [36] M.H. Cho, E.J. Lee, M. Son, J.-H. Lee, D. Yoo, J.-W. Kim, S.W. Park, J.-S. Shin, J. Cheon, A magnetic switch for the control of cell death signalling in vitro and in vivo systems, *Nat. Mater.* 11 (2012) 1038–1043.
- [37] C.W. Burke, J.S. Suk, A.J. Kim, Y.-H.J. Hsiang, A.L. Klibanov, J. Hanes, R.J. Price, Markedly enhanced skeletal muscle transfection achieved by the ultrasound-targeted delivery of non-viral gene nanocarriers with microbubbles, *J. Control. Release* 162 (2012) 414–421.
- [38] C.W. Burke, A.L. Klibanov, J.P. Sheehan, R.J. Price, Inhibition of glioma growth by microbubble activation in a subcutaneous model using low duty cycle ultrasound without significant heating, *J. Neurosurg.* 114 (2011) 1654–1661.
- [39] Y. Matsumura, H. Maeda, A new concept for macromolecular therapeutics in cancer chemotherapy: mechanism of tumorotropic accumulation of proteins and the antitumor agent smancs, *Cancer Res.* 46 (1986) 6387–6392.
- [40] H. Maeda, SMANCS and polymer-conjugated macromolecular drugs: advantages in cancer chemotherapy, *Adv. Drug Deliv. Rev.* 46 (2001) 169–185.
- [41] A. Kobayashi, T. Oda, H. Maeda, Protein binding of macromolecular anticancer agent SMANCS: characterization of poly(styrene-co-maleic acid) derivatives as an albumin binding ligand, *J. Bioact. Compat. Polym.* 3 (1988) 319–333.
- [42] D.O. Bates, N.J. Hillman, B. Williams, C.R. Neal, T.M. Pocock, Regulation of microvascular permeability by vascular endothelial growth factors*, *J. Anat.* 200 (2002) 581–597.
- [43] R.K. Jain, The next frontier of molecular medicine: delivery of therapeutics, *Nat. Med.* 4 (1998) 655–657.
- [44] R.K. Jain, T. Stylianopoulos, Delivering nanomedicine to solid tumors, *Nat. Rev. Clin. Oncol.* 7 (2010) 653–664.
- [45] S.K. Hobbs, W.L. Monsky, F. Yuan, W.G. Roberts, L. Griffith, V.P. Torchilin, R.K. Jain, Regulation of transport pathways in tumor vessels: role of tumor type and micro-environment, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 4607–4612.
- [46] M.A. Swartz, M.E. Fleury, Interstitial flow and its effects in soft tissues, *Annu. Rev. Biomed. Eng.* 9 (2007) 229–256.
- [47] T.P. Padera, B.R. Stoll, J.B. Tooredman, D. Capen, E.d. Tomaso, R.K. Jain, Pathology: cancer cells compress intratumour vessels, *Nature* 427 (2004) 695.
- [48] R.K. Jain, Transport of molecules across tumor vasculature, *Cancer Metastasis Rev.* 6 (1987) 559–593.
- [49] M.A. Swartz, The physiology of the lymphatic system, *Adv. Drug Deliv. Rev.* 50 (2001) 3–20.
- [50] Y. Noguchi, J. Wu, R. Duncan, J. Strohal, K. Ulbrich, T. Akaike, H. Maeda, Early phase tumor accumulation of macromolecules: a great difference in clearance rate between tumor and normal tissues, *Jpn. J. Cancer Res.* 89 (1998) 307–314.
- [51] J.M. Rabanel, V. Aoun, I. Elkin, M. Mokhtar, P. Hildgen, Drug-loaded nanocarriers: passive targeting and crossing of biological barriers, *Curr. Med. Chem.* 19 (2012) 3070–3102.
- [52] S.A. Stacker, M.G. Achen, L. Jussila, M.E. Baldwin, K. Alitalo, Metastasis: lymphangiogenesis and cancer metastasis, *Nat. Rev. Cancer* 2 (2002) 573–583.
- [53] T. Stylianopoulos, K. Soteriou, D. Fukumura, R.K. Jain, Cationic nanoparticles have superior transvascular flux into solid tumors: insights from a mathematical model, *Ann. Biomed. Eng.* 41 (2013) 68–77.
- [54] M. Dellian, F. Yuan, V.S. Trubetskoy, V.P. Torchilin, R.K. Jain, Vascular permeability in a human tumour xenograft: molecular charge dependence, *Br. J. Cancer* 82 (2000) 1513–1518.
- [55] M. Schmitt-Sody, S. Strieth, S. Krasnici, B. Sauer, B. Schulze, M. Teifel, U. Michaelis, K. Naujoks, M. Dellian, Neovascular targeting therapy: paclitaxel encapsulated in cationic liposomes improves antitumoral efficacy, *Clin. Cancer Res.* 9 (2003) 2335–2341.
- [56] S. Krasnici, A. Werner, M.E. Eichhorn, M. Schmitt-Sody, S.A. Pahernik, B. Sauer, B. Schulze, M. Teifel, U. Michaelis, K. Naujoks, M. Dellian, Effect of the surface charge of liposomes on their uptake by angiogenic tumor vessels, *Int. J. Cancer* 105 (2003) 561–567.
- [57] W.C. Zamboni, J.L. Eiseman, S. Strychor, P.M. Rice, E. Joseph, B.A. Zamboni, M.K. Donnelly, J. Shurer, R.A. Parise, M.E. Tonda, N.Y. Yu, P.H. Basse, Tumor disposition of pegylated liposomal CKD-602 and the reticuloendothelial system in preclinical tumor models, *J. Liposome Res.* 21 (2011) 70–80.
- [58] H. Hashizume, P. Baluk, S. Morikawa, J.W. McLean, G. Thurston, S. Roberge, R.K. Jain, D.M. McDonald, Openings between defective endothelial cells explain tumor vessel leakiness, *Am. J. Pathol.* 156 (2000) 1363–1380.
- [59] P.A. Netti, L.M. Hamberg, J.W. Babich, D. Kierstead, W. Graham, G.J. Hunter, G.L. Wolf, A. Fischman, Y. Boucher, R.K. Jain, Enhancement of fluid filtration across tumor vessels: implication for delivery of macromolecules, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 3137–3142.
- [60] O. Lieleg, R.M. Baumgärtel, A.R. Bausch, Selective filtering of particles by the extracellular matrix: an electrostatic bandpass, *Biophys. J.* 97 (2009) 1569–1577.
- [61] G. Alexandrakis, E.B. Brown, R.T. Tong, T.D. McKee, R.B. Campbell, Y. Boucher, R.K. Jain, Two-photon fluorescence correlation microscopy reveals the two-phase nature of transport in tumors, *Nat. Med.* 10 (2004) 203–207.
- [62] P.A. Netti, D.A. Berk, M.A. Swartz, A.J. Grodzinsky, R.K. Jain, Role of extracellular matrix assembly in interstitial transport in solid tumors, *Cancer Res.* 60 (2000) 2497–2503.
- [63] T.D. McKee, P. Grandi, W. Mok, G. Alexandrakis, N. Insin, J.P. Zimmer, M.G. Bawendi, Y. Boucher, X.O. Breakefield, R.K. Jain, Degradation of fibrillar collagen in a human melanoma xenograft improves the efficacy of an oncolytic Herpes Simplex virus vector, *Cancer Res.* 66 (2006) 2509–2513.
- [64] W.P. Caron, G. Song, P. Kumar, S. Rawal, W.C. Zamboni, Interpatient pharmacokinetic and pharmacodynamic variability of carrier-mediated anticancer agents, *Clin. Pharmacol. Ther.* 91 (2012) 802–812.
- [65] W.C. Zamboni, L.J. Maruca, S. Strychor, B.A. Zamboni, S. Ramalingam, R.P. Edwards, J. Kim, Y. Bang, H. Lee, D.M. Friedland, R.G. Stoller, C.P. Belani, R.K. Ramanathan, Bidirectional pharmacodynamic interaction between pegylated liposomal CKD-602 (S-CKD602) and monocytes in patients with refractory solid tumors, *J. Liposome Res.* 21 (2011) 158–165.
- [66] K. Sano, T. Nakajima, P.L. Choyke, H. Kobayashi, Markedly enhanced permeability and retention effects induced by photo-immunotherapy of tumors, *ACS Nano* 7 (2013) 717–724.
- [67] B. Diop-Frimpong, V.P. Chauhan, S. Krane, Y. Boucher, R.K. Jain, Losartan inhibits collagen I synthesis and improves the distribution and efficacy of nanotherapeutics in tumors, *Proc. Natl. Acad. Sci.* 108 (7) (2011) 2909–2914.
- [68] A. Noguchi, T. Takahashi, T. Yamaguchi, K. Kitamura, A. Noguchi, H. Tsurumi, K. Takashina, H. Maeda, Enhanced tumor localization of monoclonal antibody by treatment with kinase II inhibitor and angiotensin II, *Jpn. J. Cancer Res.* 83 (1992) 240–243.
- [69] H. Maeda, Nitroglycerin enhances vascular blood flow and drug delivery in hypoxic tumor tissues: analogy between angina pectoris and solid tumors and enhancement of the EPR effect, *J. Control. Release* 142 (2010) 296–298.
- [70] H. Maeda, Macromolecular therapeutics in cancer treatment: the EPR effect and beyond, *J. Control. Release* 164 (2012) 138–144.
- [71] J. Fang, H. Qin, H. Nakamura, K. Tsukigawa, T. Shin, H. Maeda, Carbon monoxide, generated by heme oxygenase-1, mediates the enhanced permeability and retention effect in solid tumors, *Cancer Sci.* 103 (2012) 535–541.

- [72] V.P. Chauhan, T. Stylianopoulos, J.D. Martin, Z. Popovic, O. Chen, W.S. Kamoun, M.G. Bawendi, D. Fukumura, R.K. Jain, Normalization of tumour blood vessels improves the delivery of nanomedicines in a size-dependent manner, *Nat. Nanotechnol.* 7 (2012) 384–388.
- [73] M.R. Kano, Y. Komuta, C. Iwata, M. Oka, Y.-T. Shirai, Y. Morishita, Y. Ouchi, K. Kataoka, K. Miyazono, Comparison of the effects of the kinase inhibitors imatinib, sorafenib, and transforming growth factor- β receptor inhibitor on extravasation of nanoparticles from neovasculature, *Cancer Sci.* 100 (2009) 173–180.
- [74] H. Cabral, Y. Matsumoto, K. Mizuno, Q. Chen, M. Murakami, M. Kimura, Y. Terada, M.R. Kano, K. Miyazono, M. Uesaka, N. Nishiyama, K. Kataoka, Accumulation of sub-100 nm polymeric micelles in poorly permeable tumours depends on size, *Nat. Nanotechnol.* 6 (2011) 815–823.
- [75] M.R. Kano, Y. Bae, C. Iwata, Y. Morishita, M. Yashiro, M. Oka, T. Fujii, A. Komuro, K. Kiyono, M. Kaminishi, K. Hirakawa, Y. Ouchi, N. Nishiyama, K. Kataoka, K. Miyazono, Improvement of cancer-targeting therapy, using nanocarriers for intractable solid tumors by inhibition of TGF- β signaling, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 3460–3465.
- [76] H. Maeda, The enhanced permeability and retention (EPR) effect in tumor vasculature: the key role of tumor-selective macromolecular drug targeting, *Adv. Enzyme Regul.* 41 (2001) 189–207.
- [77] M.R. Dreher, W. Liu, C.R. Michelich, M.W. Dewhirst, F. Yuan, A. Chilkoti, Tumor vascular permeability, accumulation, and penetration of macromolecular drug carriers, *JNCI, J. Natl. Cancer Inst.* 98 (2006) 335–344.
- [78] Y. Murakami, Y. Tabata, Y. Ikada, Tumor accumulation of poly(ethylene glycol) with different molecular weights after intravenous injection, *Drug Deliv.* 4 (1997) 23–31.
- [79] M.E. Fox, F.C. Szoka, J.M.J. Fréchet, Soluble polymer carriers for the treatment of cancer: the importance of the molecular architecture, *Acc. Chem. Res.* 42 (2009) 1141–1151.
- [80] T.M. Allen, M.S. Newman, M.C. Woodle, E. Mayhew, P.S. Uster, Pharmacokinetics and anti-tumor activity of vincristine encapsulated in sterically stabilized liposomes, *Int. J. Cancer* 62 (1995) 199–204.
- [81] V.P. Torchilin, Recent advances with liposomes as pharmaceutical carriers, *Nat. Rev. Drug Discov.* 4 (2005) 145–160.
- [82] S.D. Perrault, C. Walkey, T. Jennings, H.C. Fischer, W.C. Chan, Mediating tumor targeting efficiency of nanoparticles through design, *Nano Lett.* 9 (2009) 1909–1915.
- [83] S. Huo, H. Ma, K. Huang, J. Liu, T. Wei, S. Jin, J. Zhang, S. He, X.-J. Liang, Superior penetration and retention behavior of 50 nm gold nanoparticles in tumors, *Cancer Res.* 73 (2013) 319–330.
- [84] C. Wong, T. Stylianopoulos, J. Cui, J. Martin, V.P. Chauhan, W. Jiang, Z. Popović, R.K. Jain, M.G. Bawendi, D. Fukumura, Multistage nanoparticle delivery system for deep penetration into tumor tissue, *Proc. Natl. Acad. Sci. U. S. A.* 108 (2011) 2426–2431.
- [85] Y. Takakura, R.I. Mahato, M. Hashida, Extravasation of macromolecules, *Adv. Drug Deliv. Rev.* 34 (1998) 93–108.
- [86] G.L. Scherphof, J.A.A.M. Kamps, The role of hepatocytes in the clearance of liposomes from the blood circulation, *Prog. Lipid Res.* 40 (2001) 149–166.
- [87] K. Nishida, K. Mihara, T. Takino, S. Nakane, Y. Takakura, M. Hashida, H. Sezaki, Hepatic disposition characteristics of electrically charged macromolecules in rat *in vivo* and in the perfused liver, *Pharm. Res.* 8 (1991) 437–444.
- [88] C. Salvador-Morales, L. Zhang, R. Langer, O.C. Farokhzad, Immunocompatibility properties of lipid-polymer hybrid nanoparticles with heterogeneous surface functional groups, *Biomaterials* 30 (2009) 2231–2240.
- [89] A. Chonn, P.R. Cullis, D.V. Devine, The role of surface charge in the activation of the classical and alternative pathways of complement by liposomes, *J. Immunol.* 146 (1991) 4234–4241.
- [90] C. He, Y. Hu, L. Yin, C. Tang, C. Yin, Effects of particle size and surface charge on cellular uptake and biodistribution of polymeric nanoparticles, *Biomaterials* 31 (2010) 3657–3666.
- [91] K. Xiao, Y. Li, J. Luo, J.S. Lee, W. Xiao, A.M. Gonik, R.G. Agarwal, K.S. Lam, The effect of surface charge on *in vivo* biodistribution of PEG-oligocholeic acid based micellar nanoparticles, *Biomaterials* 32 (2011) 3435–3446.
- [92] T.S. Levchenko, R. Rammohan, A.N. Lukyanov, K.R. Whiteman, V.P. Torchilin, Liposome clearance in mice: the effect of a separate and combined presence of surface charge and polymer coating, *Int. J. Pharm.* 240 (2002) 95–102.
- [93] R.R. Arvizo, O.R. Miranda, D.F. Moyano, C.A. Walden, K. Giri, R. Bhattacharya, J.D. Robertson, V.M. Rotello, J.M. Reid, P. Mukherjee, Modulating pharmacokinetics, tumor uptake and biodistribution by engineered nanoparticles, *PLoS ONE* 6 (2011) e24374.
- [94] A. Gabizon, D. Papahadjopoulos, Liposome formulations with prolonged circulation time in blood and enhanced uptake by tumors, *Proc. Natl. Acad. Sci. U. S. A.* 85 (1988) 6949–6953.
- [95] E. Roux, M. Lafleur, É. Lataste, P. Moreau, J.-C. Leroux, On the characterization of pH-sensitive liposome/polymer complexes, *Biomacromolecules* 4 (2003) 240–248.
- [96] D. Peer, R. Margalit, Tumor-targeted hyaluronan nanoliposomes increase the anti-tumor activity of liposomal doxorubicin in syngeneic and human xenograft mouse tumor models, *Neoplasia* 6 (2004) 343–353.
- [97] Y. Yamamoto, Y. Nagasaki, Y. Kato, Y. Sugiyama, K. Kataoka, Long-circulating poly(ethylene glycol)-poly(D, L-lactide) block copolymer micelles with modulated surface charge, *J. Control. Release* 77 (2001) 27–38.
- [98] R.B. Campbell, D. Fukumura, E.B. Brown, L.M. Mazzola, Y. Izumi, R.K. Jain, V.P. Torchilin, L.L. Munn, Cationic charge determines the distribution of liposomes between the vascular and extravascular compartments of tumors, *Cancer Res.* 62 (2002) 6831–6836.
- [99] S. Strieth, M.E. Eichhorn, A. Werner, B. Sauer, M. Teifel, U. Michaelis, A. Berghaus, M. Dellian, Paclitaxel encapsulated in cationic liposomes increases tumor microvessel leakiness and improves therapeutic efficacy in combination with cisplatin, *Clin. Cancer Res.* 14 (2008) 4603–4611.
- [100] J.M. Löhr, S.L. Haas, W.-O. Bechstein, G. Bodoky, K. Cwiertka, W. Fischbach, U.R. Fölsch, D. Jäger, D. Osinsky, J. Prausova, W.E. Schmidt, M.P. Lutz, Cationic liposomal paclitaxel plus gemcitabine or gemcitabine alone in patients with advanced pancreatic cancer: a randomized controlled phase II trial, *Ann. Oncol.* 23 (2012) 1214–1222.
- [101] U. Fasol, A. Frost, M. Büchert, J. Arends, U. Fiedler, D. Scharr, J. Scheuenpflug, K. Mross, Vascular and pharmacokinetic effects of EndoTAG-1 in patients with advanced cancer and liver metastasis, *Ann. Oncol.* 23 (2012) 1030–1036.
- [102] H. Meng, M. Xue, T. Xia, Z. Ji, D.Y. Tam, J.J. Zink, A.E. Nel, Use of size and a copolymer design feature to improve the biodistribution and the enhanced permeability and retention effect of doxorubicin-loaded mesoporous silica nanoparticles in a murine xenograft tumor model, *ACS Nano* 5 (2011) 4131–4144.
- [103] E.A. Ho, E. Ramsay, M. Ginj, M. Anantha, I. Bregman, J. Sy, J. Woo, M. Osooly-Talesh, D.T. Yapp, M.B. Bally, Characterization of cationic liposome formulations designed to exhibit extended plasma residence times and tumor vasculature targeting properties, *J. Pharm. Sci.* 99 (2010) 2839–2853.
- [104] H.-S. Han, J.D. Martin, J. Lee, D.K. Harris, D. Fukumura, R.K. Jain, M. Bawendi, Spatial charge configuration regulates nanoparticle transport and binding behavior *in vivo*, *Angew. Chem. Int. Ed.* 52 (2013) 1414–1419.
- [105] T. Nomura, N. Koreeda, F. Yamashita, Y. Takakura, M. Hashida, Effect of particle size and charge on the disposition of lipid carriers after intratumoral injection into tissue-isolated tumors, *Pharm. Res.* 15 (1998) 128–132.
- [106] T. Nomura, A. Saikawa, S. Morita, T. Sakaeda, F. Yamashita, K. Honda, T. Yoshinobu, M. Hashida, Pharmacokinetic characteristics and therapeutic effects of mitomycin C-dextran conjugates after intratumoural injection, *J. Control. Release* 52 (1998) 239–252.
- [107] J.A. Champion, S. Mitragotri, Shape induced inhibition of phagocytosis of polymer particles, *Pharm. Res.* 26 (2009) 244–249.
- [108] J.A. Champion, S. Mitragotri, Role of target geometry in phagocytosis, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 4930–4934.
- [109] Y. Geng, P. Dalhaimer, S. Cai, R. Tsai, M. Tewari, T. Minko, D.E. Discher, Shape effects of filaments versus spherical particles in flow and drug delivery, *Nat. Nanotechnol.* 2 (2007) 249–255.
- [110] A. Ruggiero, C.H. Villa, E. Bander, D.A. Rey, M. Bergkvist, C.A. Batt, K. Manova-Todorova, W.M. Deen, D.A. Scheinberg, M.R. McDevitt, Paradoxical glomerular filtration of carbon nanotubes, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 12369–12374.
- [111] V.P. Chauhan, Z. Popović, O. Chen, J. Cui, D. Fukumura, M.G. Bawendi, R.K. Jain, Fluorescent nanorods and nanospheres for real-time *in vivo* probing of nanoparticle shape-dependent tumor penetration, *Angew. Chem. Int. Ed.* 50 (2011) 11417–11420.
- [112] S. Shukla, A.L. Abblack, A.M. Wen, K.L. Lee, J.D. Lewis, N.F. Steinmetz, Increased tumor homing and tissue penetration of the filamentous plant viral nanoparticle potato virus X, *Mol. Pharm.* 10 (2013) 33–42.
- [113] T. Lammers, F. Kiessling, W.E. Hennink, G. Storm, Drug targeting to tumors: principles, pitfalls and (pre-) clinical progress, *J. Control. Release* 161 (2012) 175–187.
- [114] S. Taurin, H. Nehoff, K. Greish, Anticancer nanomedicine and tumor vascular permeability; where is the missing link? *J. Control. Release* 164 (2012) 265–275.
- [115] Y. Liu, Y.-C. Tseng, L. Huang, Biodistribution studies of nanoparticles using fluorescence imaging: a qualitative or quantitative method? *Pharm. Res.* 29 (2012) 3273–3277.
- [116] A. Gabizon, R. Catane, B. Uziely, B. Kaufman, T. Safra, R. Cohen, F.J. Martin, A. Huang, Y. Barenholz, Prolonged circulation time and enhanced accumulation in malignant exudates of doxorubicin encapsulated in polyethylene-glycol coated liposomes, *Cancer Res.* 54 (1994) 987–992.
- [117] D.W. Northfelt, F.J. Martin, P.K. Working, P.A. Volberding, J. Russell, M. Newman, M.A. Amantea, L.D. Kaplan, Doxorubicin encapsulated in liposomes containing surface-bound polyethylene glycol: pharmacokinetics, tumor localization, and safety in patients with AIDS-related Kaposi's sarcoma, *J. Clin. Pharmacol.* 36 (1996) 55–63.
- [118] Z. Symon, A. Peyser, D. Tzemach, O. Lyass, E. Sucher, E. Shezen, A. Gabizon, Selective delivery of doxorubicin to patients with breast carcinoma metastases by stealth liposomes, *Cancer* 86 (1999) 72–78.
- [119] K.J. Harrington, S. Mohammadtaghi, P.S. Uster, D. Glass, A.M. Peters, R.G. Vile, J.S.W. Stewart, Effective targeting of solid tumors in patients with locally advanced cancers by radiolabeled PEGylated liposomes, *Clin. Cancer Res.* 7 (2001) 243–254.
- [120] M.I. Koukourakis, S. Koukouraki, A. Giatoromanolaki, S. Kakolyris, V. Georgoulas, A. Velidaki, S. Archmandritis, N. Karkavitsas, High intratumoral accumulation of stealth liposomal doxorubicin in sarcomas: rationale for combination with radiotherapy, *Acta Oncol.* 39 (2000) 207–211.
- [121] M.I. Koukourakis, S. Koukouraki, I. Fezoulidis, N. Kelekis, G. Kyrias, S. Archmandritis, N. Karkavitsas, High intratumoural accumulation of stealth liposomal doxorubicin (Caelyx(r)) in glioblastomas and in metastatic brain tumours, *Br. J. Cancer* 83 (2000) 1281–1286.
- [122] H.D. Han, A. Lee, C.K. Song, T. Hwang, H. Seong, C.O. Lee, B.C. Shin, *In vivo* distribution and antitumor activity of heparin-stabilized doxorubicin-loaded liposomes, *Int. J. Pharm.* 313 (2006) 181–188.
- [123] K.J. Harrington, G. Rowlinson-Busza, K.N. Syrigos, P.S. Uster, R.G. Vile, A.M. Peters, J.S.W. Stewart, The effect of irradiation on the biodistribution of radiolabeled pegylated liposomes, *Int. J. Radiat. Oncol. Biol. Phys.* 50 (2001) 809–820.
- [124] N. La-Beck, B. Zamboni, A. Gabizon, H. Schmeeda, M. Amantea, P. Gehrig, W. Zamboni, Factors affecting the pharmacokinetics of pegylated liposomal doxorubicin in patients, *Cancer Chemother. Pharmacol.* 69 (2012) 43–50.

- [125] A. Schroeder, D.A. Heller, M.M. Winslow, J.E. Dahlman, G.W. Pratt, R. Langer, T. Jacks, D.G. Anderson, Treating metastatic cancer with nanotechnology, *Nat. Rev. Cancer* 12 (2012) 39–50.
- [126] A. Gabizon, H. Schmeeda, Y. Barenholz, Pharmacokinetics of pegylated liposomal doxorubicin: review of animal and human studies, *Clin. Pharmacokinet.* 42 (2003) 419–436.
- [127] W. Caron, H. Clewell, R. Dedrick, R. Ramanathan, W. Davis, N. Yu, M. Tonda, J. Schellens, J. Beijnen, W. Zamboni, Allometric scaling of pegylated liposomal anticancer drugs, *J. Pharmacokinet. Pharmacodyn.* 38 (2011) 653–669.
- [128] W.C. Zamboni, V. Torchilin, A.K. Patri, J. Hrkach, S. Stern, R. Lee, A. Nel, N.J. Panaro, P. Grodzinski, Best practices in cancer nanotechnology: perspective from NCI nanotechnology alliance, *Clin. Cancer Res.* 18 (2012) 3229–3241.
- [129] S. Kummer, L. Rubinstein, R. Kinders, R.E. Parchment, M.E. Gutierrez, A.J. Murgo, J. Ji, B. Mroczkowski, O.K. Pickeral, M. Simpson, M. Hollingshead, S.X. Yang, L. Helman, R. Wiltrout, J. Collins, J.E. Tomaszewski, J.H. Doroshow, Phase 0 clinical trials: conceptions and misconceptions, *Cancer J.* 14 (2008) 133–137 (110.1097/PPO.1090b1013e318172d318176f318173).
- [130] E. Karathanasis, S. Suryanarayanan, S.R. Balusu, K. McNeeley, I. Sechopoulos, A. Karellas, A.V. Annapragada, R.V. Bellamkonda, Imaging nanoprobe for prediction of outcome of nanoparticle chemotherapy by using mammography, *Radiology* 250 (2009) 398–406.
- [131] H.E. Daldrup-Link, D. Golovko, B. Ruffell, D.G. DeNardo, R. Castaneda, C. Ansari, J. Rao, G.A. Tikhomirov, M.F. Wendland, C. Corot, L.M. Coussens, MRI of tumor-associated macrophages with clinically applicable iron oxide nanoparticles, *Clin. Cancer Res.* 17 (2011) 5695–5704.
- [132] M.G. Harisinghani, J. Barentsz, P.F. Hahn, W.M. Deserno, S. Tabatabaei, C.H. van de Kaa, J. de la Rosette, R. Weissleder, Noninvasive detection of clinically occult lymph-node metastases in prostate cancer, *N. Engl. J. Med.* 348 (2003) 2491–2499.
- [133] J.L. Gaglia, A.R. Guimaraes, M. Harisinghani, S.E. Turvey, R. Jackson, C. Benoist, D. Mathis, R. Weissleder, Noninvasive imaging of pancreatic islet inflammation in type 1A diabetes patients, *J. Clin. Invest.* 121 (2011) 442–445.
- [134] X.-Q. Zhang, X. Xu, N. Bertrand, E. Pridgen, A. Swami, O.C. Farokhzad, Interactions of nanomaterials and biological systems: implications to personalized nanomedicine, *Adv. Drug Deliv. Rev.* 64 (2012) 1363–1384.
- [135] R. Karnik, F. Gu, P. Basto, C. Cannizzaro, L. Dean, W. Kyei-Manu, R. Langer, O.C. Farokhzad, Microfluidic platform for controlled synthesis of polymeric nanoparticles, *Nano Lett.* 8 (2008) 2906–2912.
- [136] N. Kolishetti, S. Dhar, P.M. Valencia, L.Q. Lin, R. Karnik, S.J. Lippard, R. Langer, O.C. Farokhzad, Engineering of self-assembled nanoparticle platform for precisely controlled combination drug therapy, *Proc Natl Acad Sci U S A* 107 (42) (2010) 17939–1744.
- [137] S. Aryal, C.-M.J. Hu, L. Zhang, Combinatorial drug conjugation enables nanoparticle dual-drug delivery, *Small* 6 (2010) 1442–1448.
- [138] N. Bertrand, J.G. Fleischer, K.M. Wasan, J.C. Leroux, Pharmacokinetics and biodistribution of *N*-isopropylacrylamide copolymers for the design of pH-sensitive liposomes, *Biomaterials* 30 (2009) 2598–2605.
- [139] W. Gao, R. Langer, O.C. Farokhzad, Poly(ethylene glycol) with observable shedding, *Angew. Chem. Int. Ed.* 49 (2010) 6567–6571.
- [140] M.-C. Bissery, G. Nohynek, G.-J. Sanderink, F. Lavelle, Docetaxel (Taxotere®) a review of preclinical and clinical experience, Part I: preclinical experience, *Anti-Cancer Drugs* 6 (1995) 339–355.
- [141] A.V. Ullal, T. Reiner, K.S. Yang, R. Gorbатов, C. Min, D. Issadore, H. Lee, R. Weissleder, Nanoparticle-Mediated measurement of target–drug binding in cancer cells, *ACS Nano* 5 (2011) 9216–9224.
- [142] J. Shi, Z. Xiao, N. Kamaly, O.C. Farokhzad, Self-assembled targeted nanoparticles: evolution of technologies and bench to bedside translation, *Acc. Chem. Res.* 44 (2011) 1123–1134.
- [143] Z. Cheng, A. Al Zaki, J.Z. Hui, V.R. Muzykantov, A. Tsourkas, Multifunctional nanoparticles: cost versus benefit of adding targeting and imaging capabilities, *Science* 338 (2012) 903–910.
- [144] A. Koshkaryev, R. Sawant, M. Deshpande, V. Torchilin, Immunoconjugates and long circulating systems: origins, current state of the art and future directions, *Adv. Drug Deliv. Rev.* 65 (2013) 24–35.
- [145] J.D. Byrne, T. Betancourt, L. Brannon-Peppas, Active targeting schemes for nanoparticle systems in cancer therapeutics, *Adv. Drug Deliv. Rev.* 60 (2008) 1615–1626.
- [146] F. Alexis, E. Pridgen, L.K. Molnar, O.C. Farokhzad, Factors affecting the clearance and biodistribution of polymeric nanoparticles, *Mol. Pharm.* 5 (2008) 505–515.
- [147] M.P. Monopoli, C. Aberg, A. Salvati, K.A. Dawson, Biomolecular coronas provide the biological identity of nanosized materials, *Nat. Nanotechnol.* 7 (2012) 779–786.
- [148] F. Gu, L. Zhang, B.A. Tepy, N. Mann, A. Wang, A.F. Radovic-Moreno, R. Langer, O.C. Farokhzad, Precise engineering of targeted nanoparticles by using self-assembled biointegrated block copolymers, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 2586–2591.
- [149] W. Jiang, B.Y. Kim, J.T. Rutka, W.C. Chan, Nanoparticle-mediated cellular response is size-dependent, *Nat. Nanotechnol.* 3 (2008) 145–150.
- [150] P.M. Valencia, M.H. Hanewich-Hollatz, W. Gao, F. Karim, R. Langer, R. Karnik, O.C. Farokhzad, Effects of ligands with different water solubilities on self-assembly and properties of targeted nanoparticles, *Biomaterials* 32 (2011) 6226–6233.
- [151] R.N. Saha, S. Vasanthakumar, G. Bende, M. Snehaltha, Nanoparticulate drug delivery systems for cancer chemotherapy, *Mol. Membr. Biol.* 27 (2010) 215–231.
- [152] B. Yu, H.C. Tai, W. Xue, L.J. Lee, R.J. Lee, Receptor-targeted nanocarriers for therapeutic delivery to cancer, *Mol. Membr. Biol.* 27 (2010) 286–298.
- [153] A.D. Rieux, V. Pourcelle, P.D. Cani, J. Marchand-Brynaert, Y. Pr at, Targeted nanoparticles with novel non-peptidic ligands for oral delivery, *Adv. Drug Deliv. Rev.* 65 (2013) 833–844.
- [154] J. Wang, S. Tian, R.A. Petros, M.E. Napier, J. DeSimone, The complex role of multivalency in nanoparticles targeting the transferrin receptor for cancer therapies, *J. Am. Chem. Soc.* 132 (2010) 11306–11313.
- [155] J. Wu, C.-C. Chu, Water insoluble cationic poly(ester amide)s: synthesis, characterization and applications, *J. Math. Chem. B* 1 (2013) 353–360.
- [156] A.T. Florence, “Targeting” nanoparticles: the constraints of physical laws and physical barriers, *J. Control. Release* 164 (2) (2012) 115–124.
- [157] K.F. Pirolo, E.H. Chang, Does a targeting ligand influence nanoparticle tumor localization or uptake? *Trends Biotechnol.* 26 (2008) 552–558.
- [158] D.B. Kirpotin, D.C. Drummond, Y. Shao, M.R. Shalaby, K. Hong, U.B. Nielsen, J.D. Marks, C.C. Benz, J.W. Park, Antibody targeting of long-circulating lipidic nanoparticles does not increase tumor localization but does increase internalization in animal models, *Cancer Res.* 66 (2006) 6732–6740.
- [159] D.W. Bartlett, H. Su, I.J. Hildebrandt, W.A. Weber, M.E. Davis, Impact of tumor-specific targeting on the biodistribution and efficacy of siRNA nanoparticles measured by multimodality in vivo imaging, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 15549–15554.
- [160] O.C. Farokhzad, R. Langer, Impact of nanotechnology on drug delivery, *ACS Nano* 3 (2009) 16–20.
- [161] O.C. Farokhzad, J. Cheng, B.A. Tepy, I. Sherifi, S. Jon, P.W. Kantoff, J.P. Richie, R. Langer, Targeted nanoparticle–aptamer bioconjugates for cancer chemotherapy in vivo, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 6315–6320.
- [162] G. Sahay, D.Y. Alakhova, A.V. Kabanov, Endocytosis of nanomedicines, *J. Control. Release* 145 (2010) 182–195.
- [163] C. Tekle, B.V. Deurs, K. Sandvig, T.-G. Iversen, Cellular trafficking of quantum dot–ligand bioconjugates and their induction of changes in normal routing of unconjugated ligands, *Nano Lett.* 8 (2008) 1858–1865.
- [164] L.M. Bareford, P.W. Swaan, Endocytic mechanisms for targeted drug delivery, *Adv. Drug Deliv. Rev.* 59 (2007) 748–758.
- [165] S. Bhattacharyya, J.A. Khan, G.L. Curran, J.D. Robertson, R. Bhattacharya, P. Mukherjee, Efficient delivery of gold nanoparticles by dual receptor targeting, *Adv. Mater.* 23 (2011) 5034–5038.
- [166] X. Li, H. Zhou, L. Yang, G. Du, A.S. Pai–Panandiker, X. Huang, B. Yan, Enhancement of cell recognition in vitro by dual–ligand cancer targeting gold nanoparticles, *Biomaterials* 32 (2011) 2540–2545.
- [167] I.T. Papademetriou, C. Garnacho, E.H. Schuchman, S. Muro, In vivo performance of polymer nanocarriers dually-targeted to epitopes of the same or different receptors, *Biomaterials* 34 (2013) 3459–3466.
- [168] J. Gao, S.-S. Feng, Y. Guo, Antibody engineering promotes nanomedicine for cancer treatment, *Nanomedicine* 5 (2010) 1141–1145.
- [169] R. Duncan, M.J. Vicent, F. Greco, R.I. Nicholson, Polymer–drug conjugates: towards a novel approach for the treatment of endocrine-related cancer, *Endocrinol. Relat. Cancer* 12 (2005) S189–S199.
- [170] P. Ghosh, G. Han, M. De, C.K. Kim, V.M. Rotello, Gold nanoparticles in delivery applications, *Adv. Drug Deliv. Rev.* 60 (2008) 1307–1315.
- [171] W. Zhang, Z. Zhang, Y. Zhang, The application of carbon nanotubes in target drug delivery systems for cancer therapies, *Nanoscale Res. Lett.* 6 (2011) 555.
- [172] M. Liong, J. Lu, M. Kovichich, T. Xia, S.G. Ruehm, A.E. Nel, F. Tamanoi, J.I. Zink, Multifunctional inorganic nanoparticles for imaging, targeting, and drug delivery, *ACS Nano* 2 (2008) 889–896.
- [173] O. Veiseh, J.W. Gunn, M. Zhang, Design and fabrication of magnetic nanoparticles for targeted drug delivery and imaging, *Adv. Drug Deliv. Rev.* 62 (2010) 284–304.
- [174] J. Wu, D. Wu, M.A. Mutschler, C.C. Chu, Cationic hybrid hydrogels from amino-acid-based poly(ester amide): fabrication, characterization, and biological properties, *Adv. Funct. Mater.* 22 (2012) 3815–3823.
- [175] D.A. Barrera, E. Zylstra, P.T. Lansbury, R. Langer, Synthesis and RGD peptide modification of a new biodegradable copolymer: poly(lactic acid-co-lysine), *J. Am. Chem. Soc.* 115 (1993) 11010–11011.
- [176] J. Heller, J. Barr, S.Y. Ng, K.S. Abdellou, R. Gurny, Poly(ortho esters): synthesis, characterization, properties and uses, *Adv. Drug Deliv. Rev.* 54 (2002) 1015–1039.
- [177] J.R. Silvius, M.J. Zuckermann, Interbilayer transfer of phospholipid-anchored macromolecules via monomer diffusion, *Biochemistry* 32 (1993) 3153–3161.
- [178] K.E. Uhrich, S.M. Cannizzaro, R.S. Langer, K.M. Shakesheff, Polymeric systems for controlled drug release, *Chem. Rev.* 99 (1999) 3181–3198.
- [179] J.B. Delehanty, K. Boeneman, C.E. Bradburne, K. Robertson, J.E. Bongard, I.L. Medintz, Peptides for specific intracellular delivery and targeting of nanoparticles: implications for developing nanoparticle-mediated drug delivery, *Ther. Deliv.* 1 (2010) 411–433.
- [180] M. Yu, J. Park, S. Jon, Targeting strategies for multifunctional nanoparticles in cancer imaging and therapy, *Theranostics* 2 (2012) 3–44.
- [181] R. Singh, L. Kats, W.A. Bl ttler, J.M. Lambert, Formation of *N*-substituted 2-iminothiolanes when amino groups in proteins and peptides are modified by 2-iminothiolane, *Anal. Biochem.* 236 (1996) 114–125.
- [182] A. Kumar, S. Adwani, H. Dawar, G.P. Talwar, A simple method for introducing a thiol group at the 5′-end of synthetic oligonucleotides, *Nucleic Acids Res.* 19 (1991) 4561.
- [183] M. Shi, J. Lu, M.S. Shoichet, Organic nanoscale drug carriers coupled with ligands for targeted drug delivery in cancer, *J. Math. Chem.* 19 (2009) 5485–5498.
- [184] M.J. Fischer, Amine coupling through EDC/NHS: a practical approach, *Surface Plasmon, Resonance*, 2010, pp. 55–73.
- [185] J. Park, T. Mattessich, S.M. Jay, A. Agawu, W.M. Saltzman, T.M. Fahmy, Enhancement of surface ligand display on PLGA nanoparticles with amphiphilic ligand conjugates, *J. Control. Release* 156 (2011) 109–115.
- [186] T.M. Fahmy, R.M. Samstein, C.C. Harness, W. Mark Saltzman, Surface modification of biodegradable polyesters with fatty acid conjugates for improved drug targeting, *Biomaterials* 26 (2005) 5727–5736.

- [187] K. Yumura, M. Ui, H. Doi, T. Hamakubo, T. Kodama, K. Tsumoto, A. Sugiyama, Mutations for decreasing the immunogenicity and maintaining the function of core streptavidin, *Protein Sci.* 22 (2013) 213–221.
- [188] J. Rao, J. Lahiri, L. Isaacs, R.M. Weis, G.M. Whitesides, A trivalent system from vancomycin-d-Ala-d-Ala with higher affinity than avidin-biotin, *Science* 280 (1998) 708–711.
- [189] M.E. Davis, The first targeted delivery of siRNA in humans via a self-assembling, cyclodextrin polymer-based nanoparticle: from concept to clinic, *Mol. Pharm.* 6 (2009) 659–668.
- [190] D.W. Bartlett, M.E. Davis, Physicochemical and biological characterization of targeted, nucleic acid-containing nanoparticles, *Bioconjug. Chem.* 18 (2007) 456–468.
- [191] M. Mammen, S.-K. Choi, G.M. Whitesides, Polyvalent interactions in biological systems: implications for design and use of multivalent ligands and inhibitors, *Angew. Chem. Int. Ed.* 37 (1998) 2754–2794.
- [192] Y. Wu, J.A. Phillips, H. Liu, R. Yang, W. Tan, Carbon nanotubes protect DNA strands during cellular delivery, *ACS Nano* 2 (2008) 2023–2028.
- [193] D.S. Seferos, A.E. Prigodich, D.A. Giljohann, P.C. Patel, C.A. Mirkin, Polyvalent DNA nanoparticle conjugates stabilize nucleic acids, *Nano Lett.* 9 (2008) 308–311.
- [194] N. Bertrand, P. Colin, M. Ranger, J. Leblond, Designing polymeric binders for pharmaceutical applications, in: H.-J. Schneider (Ed.), *Supramolecular Systems in Biomedical Fields*, Royal Society of Chemistry, Cambridge UK, 2013, pp. 483–517.
- [195] S. Mukherjee, R.N. Ghosh, F.R. Maxfield, Endocytosis, *Physiol. Rev.* 77 (1997) 759–803.
- [196] R. Weissleder, K. Kelly, E. Yi Sun, T. Shtatland, L. Josephson, Cell-specific targeting of nanoparticles by multivalent attachment of small molecules, *Nat. Biotechnol.* 23 (2005) 1418–1423.
- [197] J.F. Stefanick, J.D. Ashley, T. Kiziltepe, B. Bilgicer, A systematic analysis of peptide linker length and liposomal polyethylene glycol coating on cellular uptake of peptide-targeted liposomes, *ACS Nano* 7 (2013) 2935–2947.
- [198] D.R. Elias, A. Poloukhina, V. Popik, A. Tsourkas, Effect of ligand density, receptor density, and nanoparticle size on cell targeting, *Nanomedicine* 9 (2013) 194–201.
- [199] Z. Poon, S. Chen, A.C. Engler, H.-I. Lee, E. Atas, G. von Maltzahn, S.N. Bhatia, P.T. Hammond, Ligand-clustered “patchy” nanoparticles for modulated cellular uptake and in vivo tumor targeting, *Angew. Chem. Int. Ed.* 49 (2010) 7266–7270.
- [200] T.M. Allen, Ligand-targeted therapeutics in anticancer therapy, *Nat. Rev. Cancer* 2 (2002) 750–763.
- [201] K. Fujimori, D.G. Covell, J.E. Fletcher, J.N. Weinstein, Modeling Analysis of the global and microscopic distribution of immunoglobulin G, F(ab')₂, and Fab in tumors, *Cancer Res.* 49 (1989) 5656–5663.
- [202] S.I. Rudnick, G.P. Adams, Affinity and avidity in antibody-based tumor targeting, *Cancer Biother. Radiopharm.* 24 (2009) 155–161.
- [203] G.M. Thurber, R. Weissleder, Quantitating antibody uptake in vivo: conditional dependence on antigen expression levels, *Mol. Imaging Biol.* 13 (2011) 623–632.
- [204] K. Fujimori, D.G. Covell, J.E. Fletcher, J.N. Weinstein, A modeling analysis of monoclonal antibody percolation through tumors: a binding-site barrier, *J. Nucl. Med.* 31 (1990) 1191–1198.
- [205] W. van Osdol, K. Fujimori, J.N. Weinstein, An analysis of monoclonal antibody distribution in microscopic tumor nodules: consequences of a “binding site barrier”, *Cancer Res.* 51 (1991) 4776–4784.
- [206] M. Juweid, R. Neumann, C. Paik, M.J. Perez-Bacete, J. Sato, W. van Osdol, J.N. Weinstein, Micropharmacology of monoclonal antibodies in solid tumors: direct experimental evidence for a binding site barrier, *Cancer Res.* 52 (1992) 5144–5153.
- [207] J.N. Weinstein, W. van Osdol, The macroscopic and microscopic pharmacology of monoclonal antibodies, *Int. J. Immunopharmacol.* 14 (1992) 457–463.
- [208] A.J. Primeau, A. Rendon, D. Hedley, L. Lilje, I.F. Tannock, The distribution of the anticancer drug doxorubicin in relation to blood vessels in solid tumors, *Clin. Cancer Res.* 11 (2005) 8782–8788.
- [209] H. Lee, H. Fong, B. Hoang, R.M. Reilly, C. Allen, The effects of particle size and molecular targeting on the intratumoral and subcellular distribution of polymeric nanoparticles, *Mol. Pharm.* 7 (2010) 1195–1208.
- [210] L. Ghitescu, M. Bendayan, Immunolabeling efficiency of protein A-gold complexes, *J. Histochem. Cytochem.* 38 (1990) 1523–1530.
- [211] S.E. Gratton, P.A. Ropp, P.D. Pohlhaus, J.C. Luft, V.J. Madden, M.E. Napier, J.M. DeSimone, The effect of particle design on cellular internalization pathways, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 11613–11618.
- [212] E. Moradi, D. Villasaliu, M. Garnett, F. Falcone, S. Stolnik, Ligand density and clustering effects on endocytosis of folate modified nanoparticles, *RSC Adv.* 2 (2012) 3025–3033.
- [213] R.A. Petros, J.M. DeSimone, Strategies in the design of nanoparticles for therapeutic applications, *Nat. Rev. Drug Discov.* 9 (2010) 615–627.
- [214] J. Wang, S. Tian, R.A. Petros, M.E. Napier, J.M. DeSimone, The complex role of multivalency in nanoparticles targeting the transferrin receptor for cancer therapies, *J. Am. Chem. Soc.* 132 (2010) 11306–11313.
- [215] S. Barua, J.-W. Yoo, P. Kolhar, A. Wakankar, Y.R. Gokarn, S. Mitragotri, Particle shape enhances specificity of antibody-displaying nanoparticles, *Proc. Natl. Acad. Sci.* 110 (9) (2013) 3270–3275.
- [216] A. Vincent, S. Babu, E. Heckert, J. Dowding, S.M. Hirst, T.M. Inerbaev, W.T. Self, C.M. Reilly, A.M.E. Masunov, T.S. Rahman, S. Seal, Protonated nanoparticle surface governing ligand tethering and cellular targeting, *ACS Nano* 3 (2009) 1203–1211.
- [217] P. Kocbek, N. Obermajer, M. Cegnar, J. Kos, J. Kristl, Targeting cancer cells using PLGA nanoparticles surface modified with monoclonal antibody, *J. Control. Release* 120 (2007) 18–26.
- [218] Y.B. Patil, U.S. Toti, A. Khadair, L. Ma, J. Panyam, Single-step surface functionalization of polymeric nanoparticles for targeted drug delivery, *Biomaterials* 30 (2009) 859–866.
- [219] F. Zhao, Y. Zhao, Y. Liu, X. Chang, C. Chen, Y. Zhao, Cellular uptake, intracellular trafficking, and cytotoxicity of nanomaterials, *Small* 7 (2011) 1322–1337.
- [220] M. Sun, L. Yang, P. Jose, L. Wang, J. Zweit, Functionalization of quantum dots with multidentate zwitterionic ligands: impact on cellular interactions and cytotoxicity, *J. Mater. Chem. B* 1 (2013) 6137–6146.
- [221] C. Hongwei, P. Hayley, I. Masayuki, S. Kanokwan, Q. Wei, C. Yong, S. Duxin, ‘Living’ PEGylation on gold nanoparticles to optimize cancer cell uptake by controlling targeting ligand and charge densities, *Nanotechnology* 24 (2013) 355101.
- [222] J. Wu, C.-C. Chu, Block copolymer of poly(ester amide) and polyesters: synthesis, characterization, and in vitro cellular response, *Acta Biomater.* 8 (2012) 4314–4323.
- [223] A. Salvati, A.S. Pitek, M.P. Monopoli, K. Prapainop, F.B. Bombelli, D.R. Hristov, P.M. Kelly, C. Aberg, E. Mahon, K.A. Dawson, Transferrin-functionalized nanoparticles lose their targeting capabilities when a biomolecule corona adsorbs on the surface, *Nat. Nanotechnol.* 8 (2013) 137–143.
- [224] A. Mori, A.M. Klibanov, V.P. Torchilin, L. Huang, Influence of the steric barrier activity of amphiphatic poly(ethylene glycol) and ganglioside GM1 on the circulation time of the liposomes and on the target binding of immunoliposomes *in vivo*, *FEBS Lett.* 284 (1991) 263–266.
- [225] S. Hak, E. Helgesen, H.H. Hektoen, E.M. Huuse, P.A. Jarzyna, W.J.M. Mulder, O. Haraldseth, C.D.L. Davies, The effect of nanoparticle polyethylene glycol surface density on ligand-directed tumor targeting studied in vivo by dual modality imaging, *ACS Nano* 6 (2012) 5648–5658.
- [226] X. Gao, Y. Cui, R.M. Levenson, L.W.K. Chung, S. Nie, In vivo cancer targeting and imaging with semiconductor quantum dots, *Nat. Biotechnol.* 22 (2004) 969–976.
- [227] T. Hamaguchi, Y. Matsumura, Y. Nakanishi, K. Muro, Y. Yamada, Y. Shimada, K. Shiroo, H. Niki, S. Hosokawa, T. Tagawa, T. Kakizoe, Antitumor effect of MCC-465, pegylated liposomal doxorubicin tagged with newly developed monoclonal antibody GAH, in colorectal cancer xenografts, *Cancer Sci.* 95 (2004) 608–613.
- [228] J.W. Park, K. Hong, D.B. Kirpotin, G. Colbern, R. Shalaby, J. Baselga, Y. Shao, U.B. Nielsen, J.D. Marks, D. Moore, D. Paphadjopoulos, C. Benz, Anti-HER2 immunoliposomes: enhanced efficacy attributable to targeted delivery, *Clin. Cancer Res.* 8 (2002) 1172–1181.
- [229] J. Winkler, P. Martin-Killias, A. Pluckthun, U. Zangemeister-Wittke, EpCAM-targeted delivery of nanocomplexed siRNA to tumor cells with designed ankyrin repeat proteins, *Mol. Cancer Ther.* 8 (2009) 2674–2683.
- [230] F. Alexis, P. Basto, E. Levy-Nissenbaum, A.F. Radovic-Moreno, L. Zhang, E. Pridgen, A.Z. Wang, S.L. Marein, K. Westerhof, L.K. Molnar, O.C. Farokhzad, HER-2-targeted nanoparticle-antibody bioconjugates for cancer therapy, *ChemMedChem* 3 (2008) 1839–1843.
- [231] P.P. Karmali, V.R. Kotamraju, M. Kastantin, M. Black, D. Missirlis, M. Tirrell, E. Ruoslahti, Targeting of albumin-embedded paclitaxel nanoparticles to tumors, *Nanomedicine* 5 (2009) 73–82.
- [232] J.-H. Park, G. von Maltzahn, L. Zhang, M.P. Schwartz, E. Ruoslahti, S.N. Bhatia, M.J. Sailor, Magnetic iron oxide nanoworms for tumor targeting and imaging, *Adv. Mater.* 20 (2008) 1630–1635.
- [233] K.N. Sugahara, T. Teesalu, P.P. Karmali, V.R. Kotamraju, L. Agemy, O.M. Girard, D. Hanahan, R.F. Mattrey, E. Ruoslahti, Tissue-penetrating delivery of compounds and nanoparticles into tumors, *Cancer Cell* 16 (2009) 510–520.
- [234] N. Graf, D.R. Bielenberg, N. Kolishetti, C. Muus, J. Banyard, O.C. Farokhzad, S.J. Lippard, $\alpha v \beta 3$ Integrin-targeted PLGA-PEG nanoparticles for enhanced anti-tumor efficacy of a Pt(IV) prodrug, *ACS Nano* 6 (2012) 4530–4539.
- [235] N. Kamaly, G. Fredman, M. Subramanian, S. Gadde, A. Pestic, L. Cheung, Z.A. Fayad, R. Langer, I. Tabas, O.C. Farokhzad, Development and in vivo efficacy of targeted polymeric inflammation-resolving nanoparticles, *Proc Natl Acad Sci U S A* 110 (16) (2013) 6506–6511.
- [236] J.M. Chan, L. Zhang, R. Tong, D. Ghosh, W. Gao, G. Liao, K.P. Yuet, D. Gray, J.-W. Rhee, J. Cheng, G. Golomb, P. Libby, R. Langer, O.C. Farokhzad, Spatiotemporal controlled delivery of nanoparticles to injured vasculature, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 2213–2218.
- [237] J.M. Chan, J.-W. Rhee, C.L. Drum, R.T. Bronson, G. Golomb, R. Langer, O.C. Farokhzad, In vivo prevention of arterial restenosis with paclitaxel-encapsulated targeted lipid-polymeric nanoparticles, *Proc. Natl. Acad. Sci.* 108 (2011) 19347–19352.
- [238] P.E. Saw, S. Kim, I.-H. Lee, J. Park, M. Yu, J. Lee, J.-I. Kim, S. Jon, Aptide-conjugated liposome targeting tumor-associated fibronectin for glioma therapy, *J. Math. Chem.* B 1 (2013) 4723–4726.
- [239] J. Cheng, B.A. Teplý, I. Sherifi, J. Sung, G. Luther, F.X. Gu, E. Levy-Nissenbaum, A.F. Radovic-Moreno, R. Langer, O.C. Farokhzad, Formulation of functionalized PLGA-PEG nanoparticles for in vivo targeted drug delivery, *Biomaterials* 28 (2007) 869–876.
- [240] D. Kim, Y.Y. Jeong, S. Jon, A drug-loaded aptamer-gold nanoparticle bioconjugate for combined ct imaging and therapy of prostate cancer, *ACS Nano* 4 (2010) 3689–3696.
- [241] Z. Xiao, C. Ji, J. Shi, E.M. Pridgen, J. Frieder, J. Wu, O.C. Farokhzad, DNA self-assembly of targeted near-infrared-responsive gold nanoparticles for cancer thermochemotherapy, *Angew. Chem. Int. Ed.* 51 (2012) 11853–11857.
- [242] M.E. Werner, S. Karve, R. Sukumar, N.D. Cummings, J.A. Copp, R.C. Chen, T. Zhang, A.Z. Wang, Folate-targeted nanoparticle delivery of chemo- and radiotherapeutics for the treatment of ovarian cancer peritoneal metastasis, *Biomaterials* 32 (2011) 8548–8554.
- [243] S. Marrache, S. Dhar, Engineering of blended nanoparticle platform for delivery of mitochondria-acting therapeutics, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 16288–16293.
- [244] D.B. Kirpotin, D.C. Drummond, Y. Shao, M.R. Shalaby, K. Hong, U.B. Nielsen, J.D. Marks, C.C. Benz, J.W. Park, Antibody targeting of long-circulating lipidic

- nanoparticles does not increase tumor localization but does increase internalization in animal models, *Cancer Res.* 66 (2006) 6732–6740.
- [245] F.R. Brennan, L. Shaw, M.G. Wing, C. Robinson, Preclinical safety testing of biotechnology-derived pharmaceuticals – understanding the issues and addressing the challenges, *Mol. Biotechnol.* 27 (2004) 59–74.
- [246] W.C. Weinberg, M.R. Frazier-Jessen, W.J. Wu, A. Weir, M. Hartsough, P. Keegan, C. Fuchs, Development and regulation of monoclonal antibody products: challenges and opportunities, *Cancer Metastasis Rev.* 24 (2005) 569–584.
- [247] P. Simard, J.C. Leroux, *In vivo* evaluation of pH-sensitive polymer-based immunoliposomes targeting the CD33 antigen, *Mol. Pharm.* 7 (2010) 1098–1107.
- [248] S.M. Ansell, P.G. Tardi, S.S. Buchkowsky, 3-(2-Pyridylidithio)propionic acid hydrazide as a cross-linker in the formation of liposome–antibody conjugates, *Bioconjug. Chem.* 7 (1996) 490–496.
- [249] G.A. Koning, H.W.M. Morselt, A. Gorter, T.M. Allen, S. Zalipsky, J.A.A.M. Kamps, G.L. Scherphof, Pharmacokinetics of differently designed immunoliposome formulations in rats with or without hepatic colon cancer metastases, *Pharm. Res.* 18 (2001) 1291–1298.
- [250] K. Jain, Advances in the field of nanooncology, *BMC Med.* 8 (2010) 83.
- [251] J.M. Nemunaitis, J. Nemunaitis, Potential of Advexin®: a p53 gene-replacement therapy in Li–Fraumeni syndrome, *Future Oncol.* 4 (2008) 759–768.
- [252] J. Nemunaitis, N. Senzer, C. Bedell, R. Nunan, L. Sleer, E. Chang, A phase I study of escalating doses of SGT-53 for intravenous infusion of patients with advanced solid tumors, *Mol. Ther.* 17 (2009) S226.
- [253] C. Mamot, R. Ritschard, A. Wicki, G. Stehle, T. Dieterle, L. Bubendorf, C. Hilker, S. Deuster, R. Herrmann, C. Rochlitz, Tolerability, safety, pharmacokinetics, and efficacy of doxorubicin-loaded anti-EGFR immunoliposomes in advanced solid tumours: a phase 1 dose-escalation study, *Lancet Oncol.* 13 (2012) 1234–1241.
- [254] L.Y.T. Chou, K. Ming, W.C.W. Chan, Strategies for the intracellular delivery of nanoparticles, *Chem. Soc. Rev.* 40 (2011) 233–245.
- [255] C.H.J. Choi, C.A. Alabi, P. Webster, M.E. Davis, Mechanism of active targeting in solid tumors with transferrin-containing gold nanoparticles, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 1235–1240.
- [256] S.K. Sahoo, W. Ma, V. Labhasetwar, Efficacy of transferrin-conjugated paclitaxel-loaded nanoparticles in a murine model of prostate cancer, *Int. J. Cancer* 112 (2004) 335–340.
- [257] P. Colas, The eleven-year switch of peptide aptamers, *J. Biol.* 7 (2008) 2.
- [258] L. Meng, J.H. Wong, L.J. Feldman, P.G. Lemaux, B.B. Buchanan, A membrane-associated thioredoxin required for plant growth moves from cell to cell, suggestive of a role in intercellular communication, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 3900–3905.
- [259] Z. Xiao, O.C. Farokhzad, Aptamer-functionalized nanoparticles for medical applications: challenges and opportunities, *ACS Nano* 6 (2012) 3670–3676.
- [260] A. Bendele, J. Seely, C. Richey, G. Sennello, G. Shopp, Renal tubular vacuolation in animals treated with polyethylene-glycol-conjugated proteins, *Toxicol. Sci.* 42 (1998) 152–157.
- [261] R. Webster, É. Didier, P. Harris, N. Siegel, J. Stadler, L. Tilbury, D. Smith, PEGylated proteins: evaluation of their safety in the absence of definitive metabolism studies, *Drug Metab. Dispos.* 35 (2007) 9–16.
- [262] P.A. Burke, S.J. DeNardo, L.A. Miers, K.R. Lamborn, S. Matzku, G.L. DeNardo, Cilengitide targeting of $\alpha v \beta 3$ integrin receptor synergizes with radioimmunotherapy to increase efficacy and apoptosis in breast cancer xenografts, *Cancer Res.* 62 (2002) 4263–4272.
- [263] S.L. Goodman, G. Hölzemann, G.A.G. Sulyok, H. Kessler, Nanomolar small molecule inhibitors for $\alpha v \beta 6$, $\alpha v \beta 5$, and $\alpha v \beta 3$ integrins, *J. Med. Chem.* 45 (2002) 1045–1051.
- [264] G. Colombo, F. Curnis, G.M.S. De Mori, A. Gasparri, C. Longoni, A. Sacchi, R. Longhi, A. Corti, Structure–activity relationships of linear and cyclic peptides containing the NGR tumor-homing motif, *J. Biol. Chem.* 277 (2002) 47891–47897.
- [265] P. Laakkonen, K. Porkka, J.A. Hoffman, E. Ruoslahti, A tumor-homing peptide with a targeting specificity related to lymphatic vessels, *Nat. Med.* 8 (2002) 751–755.
- [266] K. Porkka, P. Laakkonen, J.A. Hoffman, M. Bernasconi, E. Ruoslahti, A fragment of the HMGN2 protein homes to the nuclei of tumor cells and tumor endothelial cells *in vivo*, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 7444–7449.
- [267] L. Roth, L. Agemy, V.R. Kotamraju, G. Braun, T. Teesalu, K.N. Sugahara, J. Hamzah, E. Ruoslahti, Transmembrane targeting enabled by a novel neuropilin-binding peptide, *Oncogene* 31 (2012) 3754–3763.
- [268] G. Luo, X. Yu, C. Jin, F. Yang, D. Fu, J. Long, J. Xu, C. Zhan, W. Lu, LyP-1-conjugated nanoparticles for targeting drug delivery to lymphatic metastatic tumors, *Int. J. Pharm.* 385 (2010) 150–156.
- [269] K.N. Sugahara, T. Teesalu, P.P. Karmali, V.R. Kotamraju, L. Agemy, D.R. Greenwald, E. Ruoslahti, Coadministration of a tumor-penetrating peptide enhances the efficacy of cancer drugs, *Science* 328 (2010) 1031–1035.
- [270] S.F. Slovin, Targeting novel antigens for prostate cancer treatment: focus on prostate-specific membrane antigen, *Expert Opin. Ther. Targets* 9 (2005) 561–570.
- [271] R. Pasqualini, E. Koivunen, R. Kain, J. Lahdenranta, M. Sakamoto, A. Stryhn, R.A. Ashmun, L.H. Shapiro, W. Arap, E. Ruoslahti, Aminopeptidase N is a receptor for tumor-homing peptides and a target for inhibiting angiogenesis, *Cancer Res.* 60 (2000) 722–727.
- [272] N. Zhang, C. Chittasupho, C. Duangrat, T.J. Siahaan, C. Berkland, PLGA nanoparticle-peptide conjugate effectively targets intercellular cell-adhesion molecule-1, *Bioconjug. Chem.* 19 (2008) 145–152.
- [273] W. Hild, K. Pollinger, A. Caporale, C. Cabrele, M. Keller, N. Pluym, A. Buschauer, R. Rachel, J. Tessmar, M. Breunig, A. Goepferich, G protein-coupled receptors function as logic gates for nanoparticle binding and cell uptake, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 10667–10672.
- [274] V.B. Pinheiro, A.I. Taylor, C. Cozens, M. Abramov, M. Renders, S. Zhang, J.C. Chaput, J. Wengel, S.-Y. Peak-Chew, S.H. McLaughlin, P. Herdewijn, P. Holliger, Synthetic genetic polymers capable of heredity and evolution, *Science* 336 (2012) 341–344.
- [275] C.-L. Zhu, C.-H. Lu, X.-Y. Song, H.-H. Yang, X.-R. Wang, Bioresponsive controlled release using mesoporous silica nanoparticles capped with aptamer-based molecular gate, *J. Am. Chem. Soc.* 133 (2011) 1278–1281.
- [276] J. Lapointe, T.Q. Truong, L. Falstraalt, L. Brissette, Differential abilities of mouse liver parenchymal and nonparenchymal cells in HDL and LDL (native and oxidized) association and cholesterol efflux, *Biochem. Cell Biol.* 84 (2006) 250–256.
- [277] A. Nahvi, N. Sudarsan, M.S. Ebert, X. Zou, K.L. Brown, R.R. Breaker, Genetic control by a metabolite binding mRNA, *Chem. Biol.* 9 (2002) 1043–1049.
- [278] A.S. Mironov, I. Gusarov, R. Rafikov, L.E. Lopez, K. Shatalin, R.A. Kreneva, D.A. Perumov, E. Nudler, Sensing small molecules by nascent RNA: a mechanism to control transcription in bacteria, *Cell* 111 (2002) 747–756.
- [279] W. Winkler, A. Nahvi, R.R. Breaker, Thiamine derivatives bind messenger RNAs directly to regulate bacterial gene expression, *Nature* 419 (2002) 952–956.
- [280] A.R. Oliphant, C.J. Brandl, K. Struhl, Defining the sequence specificity of DNA-binding proteins by selecting binding sites from random-sequence oligonucleotides: analysis of yeast GCN4 protein, *Mol. Cell. Biol.* 9 (1989) 2944–2949.
- [281] C. Tuerk, L. Gold, Systematic evolution of ligands by exponential enrichment – RNA ligands to bacteriophage-T4 DNA-polymerase, *Science* 249 (1990) 505–510.
- [282] A.D. Ellington, J.W. Szostak, *In vitro* selection of RNA molecules that bind specific ligands, *Nature* 346 (1990) 818–822.
- [283] D.A. Daniels, H. Chen, B.J. Hicke, K.M. Swiderek, L. Gold, A tenascin-C aptamer identified by tumor cell SELEX: systematic evolution of ligands by exponential enrichment, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 15416–15421.
- [284] Z. Xiao, E. Levy-Nissenbaum, F. Alexis, A. Luptak, B.A. Teply, J.M. Chan, J. Shi, E. Digga, J. Cheng, R. Langer, O.C. Farokhzad, Engineering of targeted nanoparticles for cancer therapy using internalizing aptamers isolated by cell-uptake selection, *ACS Nano* 6 (2012) 696–704.
- [285] J. Mi, Y. Liu, Z.N. Rabbani, Z. Yang, J.H. Urban, B.A. Sullenger, B.M. Clary, *In vivo* selection of tumor-targeting RNA motifs, *Nat. Chem. Biol.* 6 (2010) 22–24.
- [286] O.C. Farokhzad, S. Jon, A. Khademhosseini, T.-N.T. Tran, D.A. LaVan, R. Langer, Nanoparticle–aptamer bioconjugates: a new approach for targeting prostate cancer cells, *Cancer Res.* 64 (2004) 7668–7672.
- [287] V. Bagalkot, O.C. Farokhzad, R. Langer, S. Jon, An aptamer–doxorubicin physical conjugate as a novel targeted drug-delivery platform, *Angew. Chem. Int. Ed.* 45 (2006) 8149–8152.
- [288] O.C. Farokhzad, R. Langer, Nanomedicine: developing smarter therapeutic and diagnostic modalities, *Adv. Drug Deliv. Rev.* 58 (2006) 1456–1459.
- [289] A.D. Keefe, S. Pai, A. Ellington, Aptamers as therapeutics, *Nat. Rev. Drug Discov.* 9 (2010) 537–550.
- [290] J.F. Lee, G.M. Stovall, A.D. Ellington, Aptamer therapeutics advance, *Curr. Opin. Chem. Biol.* 10 (2006) 282–289.
- [291] R.E. Kleiner, C.E. Dumelin, G.C. Tiu, K. Sakurai, D.R. Liu, *In vitro* selection of a DNA-templated small-molecule library reveals a class of macrocyclic kinase inhibitors, *J. Am. Chem. Soc.* 132 (2010) 11779–11791.
- [292] S. Melkko, J. Scheuermann, C.E. Dumelin, D. Neri, Encoded self-assembling chemical libraries, *Nat. Biotechnol.* 22 (2004) 568–574.
- [293] A.R. Hilgenbrink, P.S. Low, Folate receptor-mediated drug targeting: from therapeutics to diagnostics, *J. Pharm. Sci.* 94 (2005) 2135–2146.
- [294] S. Markert, S. Lassmann, B. Gabriel, M. Klar, M. Werner, G. Gitsch, F. Kratz, A. Hasenburger, Alpha-folate receptor expression in epithelial ovarian carcinoma and non-neoplastic ovarian tissue, *Anticancer Res.* 28 (2008) 3567–3572.
- [295] G.M. van Dam, G. Themelis, L.M.A. Crane, N.J. Harlaar, R.G. Pleijhuis, W. Kelder, A. Sarantopoulos, J.S. de Jong, H.J.G. Arts, A.G.J. van der Zee, J. Bart, P.S. Low, V. Ntziachristos, Intraoperative tumor-specific fluorescence imaging in ovarian cancer by folate receptor- α targeting: first in-human results, *Nat. Med.* 17 (2011) 1315–1319.
- [296] R.E. Fisher, B.A. Siegel, S.L. Edell, N.M. Oyesiku, D.E. Morgenstern, R.A. Messmann, R.J. Amato, Exploratory study of ^{99m}Tc -EC20 imaging for identifying patients with folate receptor-positive solid tumors, *J. Nucl. Med.* 49 (2008) 899–906.
- [297] N. Parker, M.J. Turk, E. Westrick, J.D. Lewis, P.S. Low, C.P. Leamon, Folate receptor expression in carcinomas and normal tissues determined by a quantitative radioligand binding assay, *Anal. Biochem.* 338 (2005) 284–293.
- [298] B.A. Kamen, M.T. Wang, A.J. Streckfuss, X. Peryea, R.G. Anderson, Delivery of folates to the cytoplasm of MA104 cells is mediated by a surface membrane receptor that recycles, *J. Biol. Chem.* 263 (1988) 13602–13609.
- [299] R.A.J. Smith, C.M. Porteous, A.M. Gane, M.P. Murphy, Delivery of bioactive molecules to mitochondria *in vivo*, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 5407–5412.
- [300] H. Zhang, Y. Ma, X.-L. Sun, Recent developments in carbohydrate-decorated targeted drug/gene delivery, *Med. Res. Rev.* 30 (2010) 270–289.
- [301] M. Hashida, M. Nishikawa, F. Yamashita, Y. Takakura, Cell-specific delivery of genes with glycosylated carriers, *Adv. Drug Deliv. Rev.* 52 (2001) 187–196.
- [302] J.M. Bergen, H.A. Von Recum, T.T. Goodman, A.P. Massey, S.H. Pun, Gold nanoparticles as a versatile platform for optimizing physicochemical parameters for targeted drug delivery, *Macromol. Biosci.* 6 (2006) 506–516.
- [303] D.E. Leckband, Novel functions and binding mechanisms of carbohydrate-binding proteins determined by force measurements, *Curr. Protein Pept. Sci.* 12 (2011) 743–759.
- [304] S. André, B. Frisch, H. Kaltner, D. Desouza, F. Schuber, H.-J. Gabius, Lectin-mediated drug targeting: selection of valency, sugar type (Gal/Lac), and spacer length for cluster glycosides as parameters to distinguish ligand binding to C-type asialoglycoprotein receptors and galectins, *Pharm. Res.* 17 (2000) 985–990.
- [305] L.W. Seymour, D.R. Ferry, D. Anderson, S. Hesselwood, P.J. Julian, R. Poyner, J. Doran, A.M. Young, S. Burtles, D.J. Kerr, Hepatic drug targeting: phase I evaluation of polymer-bound doxorubicin, *J. Clin. Oncol.* 20 (2002) 1668–1676.

- [306] R. Duncan, Polymer conjugates as anticancer nanomedicines, *Nat. Rev. Cancer* 6 (2006) 688–701.
- [307] S.M. Hillier, K.P. Maresca, F.J. Femia, J.C. Marquis, C.A. Foss, N. Nguyen, C.N. Zimmerman, J.A. Barrett, W.C. Eckelman, M.G. Pomper, J.L. Joyal, J.W. Babich, Preclinical evaluation of novel glutamate–urea–lysine analogues that target prostate-specific membrane antigen as molecular imaging pharmaceuticals for prostate cancer, *Cancer Res.* 69 (2009) 6932–6940.
- [308] [www.clinicaltrials.gov\(NCT00355888, in\)](http://www.clinicaltrials.gov(NCT00355888, in)).
- [309] Y. Matsumura, M. Gotoh, K. Muro, Y. Yamada, K. Shirao, Y. Shimada, M. Okuwa, S. Matsumoto, Y. Miyata, H. Ohkura, K. Chin, S. Baba, T. Yamao, A. Kannami, Y. Takamatsu, K. Ito, K. Takahashi, Phase I and pharmacokinetic study of MCC-465, a doxorubicin (DXR) encapsulated in PEG immunoliposome, in patients with metastatic stomach cancer, *Ann. Oncol.* 15 (2004) 517–525.
- [310] N. Senzer, J. Nemunaitis, D. Nemunaitis, C. Bedell, G. Edelman, M. Barve, R. Nunan, K.F. Pirolo, A. Rait, E.H. Chang, Phase I study of a systemically delivered p53 nanoparticle in advanced solid tumors, *Mol. Ther.* 21 (2013) 1096–1103.
- [311] [www.clinicaltrials.gov\(NCT01304797, in\)](http://www.clinicaltrials.gov(NCT01304797, in)).
- [312] T. Wickham, K. Futch, A phase I study of MM-302, a HER2-targeted liposomal doxorubicin, Patients with Advanced, HER2-Positive Breast Cancer, Suppl. 3, Cancer Research, San Antonio, TX, 2012, (P5-18-09).
- [313] S.P. Chawla, V.S. Chua, L. Fernandez, D. Quon, W.C. Blackwelder, E.M. Gordon, F.L. Hall, Advanced phase I/II studies of targeted gene delivery in vivo: intravenous rexin-G for gemcitabine-resistant metastatic pancreatic cancer, *Mol. Ther.* 18 (2010) 435–441.
- [314] S.P. Chawla, V.S. Chua, L. Fernandez, D. Quon, A. Saralou, W.C. Blackwelder, F.L. Hall, E.M. Gordon, Phase I/II and phase II studies of targeted gene delivery in vivo: intravenous rexin-G for chemotherapy-resistant sarcoma and osteosarcoma, *Mol. Ther.* 17 (2009) 1651–1657.
- [315] E. Galanis, S.K. Carlson, N.R. Foster, V. Lowe, F. Quevedo, R.R. McWilliams, A. Grothey, A. Jatoi, S.R. Alberts, J. Rubin, Phase I Trial of a Pathotropic Retroviral Vector Expressing a Cytocidal Cyclin G1 Construct (Rexin-G) in Patients With Advanced Pancreatic Cancer, *Mol. Ther.* 16 (2008) 979–984.
- [316] R. van der Meel, L.J.C. Vehmeijer, R.J. Kok, G. Storm, E.V.B. van Gaal, Ligand-targeted particulate nanomedicines undergoing clinical evaluation: current status, *Adv. Drug Deliv. Rev.* 65 (2013) 1284–1298.
- [317] C. Rochlitz, R. Ritschard, B. Vogel, T. Dieterle, L. Bubendorf, C. Hilker, S. Deuster, R. Herrmann, C. Mamot, A phase I study of doxorubicin-loaded anti-EGFR immunoliposomes in patients with advanced solid tumors, *Onkologie* 34 (2011) 109–109.
- [318] P. Burgess, P.B. Hutt, O.C. Farokhzad, R. Langer, S. Minick, S. Zale, On firm ground: IP protection of therapeutic nanoparticles, *Nat. Biotechnol.* 28 (2010) 1267–1270.
- [319] T.R. Daniels, T. Delgado, J.A. Rodriguez, G. Helguera, M.L. Penichet, The transferrin receptor part I: biology and targeting with cytotoxic antibodies for the treatment of cancer, *Clin. Immunol.* 121 (2006) 144–158.
- [320] A.V. Yurkovetskiy, R.J. Fram, XMT-1001, a novel polymeric camptothecin pro-drug in clinical development for patients with advanced cancer, *Adv. Drug Deliv. Rev.* 61 (2009) 1193–1202.
- [321] S. Danhauser-Riedl, E. Hausmann, H.-D. Schick, R. Bender, H. Dietschfelbinger, J. Rastetter, A.-R. Hanauske, Phase I clinical and pharmacokinetic trial of dextran conjugated doxorubicin (AD-70, DOX-OXD), *Invest. New Drugs* 11 (1993) 187–195.
- [322] N.E. Schoemaker, C. van Kesteren, H. Rosing, S. Jansen, M. Swart, J. Lieverst, D. Fraier, M. Breda, C. Pellizzoni, R. Spinelli, M.G. Porro, J.H. Beijnen, J.H.M. Schellens, W.W. ten Bokkel Huinink, A phase I and pharmacokinetic study of MAG-CPT, a water-soluble polymer conjugate of camptothecin, *Br. J. Cancer* 87 (2002) 608–614.
- [323] D. Lorusso, A. Di Stefano, V. Carone, A. Fagotti, S. Pisconti, G. Scambia, Pegylated liposomal doxorubicin-related palmar-plantar erythrodysesthesia ('hand-foot' syndrome), *Ann. Oncol.* 18 (2007) 1159–1164.
- [324] J. Szebeni, F. Muggia, A. Gabizon, Y. Barenholz, Activation of complement by therapeutic liposomes and other lipid excipient-based therapeutic products: prediction and prevention, *Adv. Drug Deliv. Rev.* 63 (2011) 1020–1030.
- [325] A.K. Rajasekaran, G. Anilkumar, J.J. Christiansen, Is prostate-specific membrane antigen a multifunctional protein? *Am. J. Physiol. Cell Physiol.* 288 (2005) C975–C981.
- [326] F.S. Ong, K. Das, J. Wang, H. Vakili, J.Z. Kuo, W.-L.B. Blackwell, S.W. Lim, M.O. Goodarzi, K.E. Bernstein, J.I. Rotter, W.W. Grody, Personalized medicine and pharmacogenetic biomarkers: progress in molecular oncology testing, *Expert. Rev. Mol. Diagn.* 12 (2012) 593–602.
- [327] N.P. Desai, V. Trieu, B. Damascelli, P. Soon-Shiong, SPARC expression correlates with tumor response to albumin-bound paclitaxel in head and neck cancer patients, *Transl. Oncol.* 2 (2009) 59–64.